

表 500 透析前補正カルシウム濃度(mg/dL) 透析前リン濃度(mg/dL)別 (透析患者全体 インタクトPTH値60～240)

透析前リン濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
3未満	36	15	78	255	91	1,055	1,508	976	145	346	199	84	44	19	4,851	41	4,892	9.29	0.83
(%)	(0.7)	(0.3)	(1.6)	(5.3)	(1.9)	(21.7)	(31.1)	(20.1)	(3.0)	(7.1)	(4.1)	(1.7)	(0.9)	(0.4)	(100.0)				
3.0～	22	20	94	422	147	1,769	2,177	1,271	156	358	193	58	24	22	6,733	47	6,780	9.20	0.72
(%)	(0.3)	(0.3)	(1.4)	(6.3)	(2.2)	(26.3)	(32.3)	(18.9)	(2.3)	(5.3)	(2.9)	(0.9)	(0.4)	(0.3)	(100.0)				
3.5～	31	47	199	742	243	3,170	4,142	2,322	262	569	265	73	27	24	12,116	85	12,201	9.17	0.66
(%)	(0.3)	(0.4)	(1.6)	(6.1)	(2.0)	(26.2)	(34.2)	(19.2)	(2.2)	(4.7)	(2.2)	(0.6)	(0.2)	(0.2)	(100.0)				
4.0～	57	53	284	1,216	404	4,985	6,158	3,494	349	839	377	109	41	37	18,403	127	18,530	9.15	0.67
(%)	(0.3)	(0.3)	(1.5)	(6.6)	(2.2)	(27.1)	(33.5)	(19.0)	(1.9)	(4.6)	(2.0)	(0.6)	(0.2)	(0.2)	(100.0)				
4.5～	55	70	379	1,489	444	5,958	7,347	4,099	458	1,086	470	129	43	27	22,054	124	22,178	9.15	0.66
(%)	(0.2)	(0.3)	(1.7)	(6.8)	(2.0)	(27.0)	(33.3)	(18.6)	(2.1)	(4.9)	(2.1)	(0.6)	(0.2)	(0.1)	(100.0)				
5.0～	76	89	391	1,634	504	5,924	7,337	4,381	454	1,112	513	117	39	19	22,590	162	22,752	9.15	0.67
(%)	(0.3)	(0.4)	(1.7)	(7.2)	(2.2)	(26.2)	(32.5)	(19.4)	(2.0)	(4.9)	(2.3)	(0.5)	(0.2)	(0.1)	(100.0)				
5.5～	63	84	378	1,455	439	5,059	6,120	3,762	423	1,075	430	123	40	25	19,476	101	19,577	9.15	0.68
(%)	(0.3)	(0.4)	(1.9)	(7.5)	(2.3)	(26.0)	(31.4)	(19.3)	(2.2)	(5.5)	(2.2)	(0.6)	(0.2)	(0.1)	(100.0)				
6.0～	10	14	82	268	66	820	1,089	639	72	164	62	18	8	3	3,315	16	3,331	9.13	0.70
(%)	(0.3)	(0.4)	(2.5)	(8.1)	(2.0)	(24.7)	(32.9)	(19.3)	(2.2)	(4.9)	(1.9)	(0.5)	(0.2)	(0.1)	(100.0)				
6.1～	38	60	237	934	274	2,832	3,223	2,096	288	602	246	81	21	13	10,945	85	11,030	9.14	0.70
(%)	(0.3)	(0.5)	(2.2)	(8.5)	(2.5)	(25.9)	(29.4)	(19.2)	(2.6)	(5.5)	(2.2)	(0.7)	(0.2)	(0.1)	(100.0)				
6.5～	52	52	230	775	257	2,409	2,739	1,767	210	515	227	65	25	12	9,335	62	9,397	9.13	0.73
(%)	(0.6)	(0.6)	(2.5)	(8.3)	(2.8)	(25.8)	(29.3)	(18.9)	(2.2)	(5.5)	(2.4)	(0.7)	(0.3)	(0.1)	(100.0)				
7.0～	35	48	168	535	134	1,378	1,611	1,036	138	309	143	54	18	12	5,619	33	5,652	9.12	0.76
(%)	(0.6)	(0.9)	(3.0)	(9.5)	(2.4)	(24.5)	(28.7)	(18.4)	(2.5)	(5.5)	(2.5)	(1.0)	(0.3)	(0.2)	(100.0)				
7.5～	20	33	120	315	75	785	924	574	75	192	106	32	5	4	3,260	34	3,294	9.11	0.77
(%)	(0.6)	(1.0)	(3.7)	(9.7)	(2.3)	(24.1)	(28.3)	(17.6)	(2.3)	(5.9)	(3.3)	(1.0)	(0.2)	(0.1)	(100.0)				
8.0～	19	11	64	198	60	451	551	316	26	113	36	12	7	6	1,870	21	1,891	9.07	0.80
(%)	(1.0)	(0.6)	(3.4)	(10.6)	(3.2)	(24.1)	(29.5)	(16.9)	(1.4)	(6.0)	(1.9)	(0.6)	(0.4)	(0.3)	(100.0)				
8.5～	12	15	53	123	26	268	311	192	22	51	23	11	6	1	1,114	14	1,128	9.03	0.82
(%)	(1.1)	(1.3)	(4.8)	(11.0)	(2.3)	(24.1)	(27.9)	(17.2)	(2.0)	(4.6)	(2.1)	(1.0)	(0.5)	(0.1)	(100.0)				
9.0～	28	28	87	213	53	372	431	254	30	92	41	12	3	2	1,646	26	1,672	8.96	0.87
(%)	(1.7)	(1.7)	(5.3)	(12.9)	(3.2)	(22.6)	(26.2)	(15.4)	(1.8)	(5.6)	(2.5)	(0.7)	(0.2)	(0.1)	(100.0)				
合計	554	639	2,844	10,574	3,217	37,235	45,668	27,179	3,108	7,423	3,331	978	351	226	143,327	978	144,305	9.15	0.70
(%)	(0.4)	(0.4)	(2.0)	(7.4)	(2.2)	(26.0)	(31.9)	(19.0)	(2.2)	(5.2)	(2.3)	(0.7)	(0.2)	(0.2)	(100.0)				
記載なし			4	9	1	16	20	20	6	3	1	2	1	1	84	152	236	9.25	0.86
(%)			(4.8)	(10.7)	(1.2)	(19.0)	(23.8)	(23.8)	(7.1)	(3.6)	(1.2)	(2.4)	(1.2)	(1.2)	(100.0)				
総計	554	639	2,848	10,583	3,218	37,251	45,688	27,199	3,114	7,426	3,332	980	352	227	143,411	1,130	144,541	9.15	0.70
(%)	(0.4)	(0.4)	(2.0)	(7.4)	(2.2)	(26.0)	(31.9)	(19.0)	(2.2)	(5.2)	(2.3)	(0.7)	(0.2)	(0.2)	(100.0)				
平均	5.63	5.76	5.59	5.42	5.32	5.18	5.14	5.17	5.19	5.23	5.15	5.14	4.94	4.86	5.20	5.32	5.20		
標準偏差	1.84	1.72	1.59	1.44	1.41	1.31	1.32	1.34	1.38	1.40	1.47	1.55	1.67	1.59	1.36	1.58	1.36		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査