

表 504 性別 原疾患別（透析患者全体 75歳以上）

原疾患	男性	女性	合計	記載なし	総計
慢性糸球体腎炎	14,780	11,809	26,589		26,589
(%)	(24.9)	(28.8)	(26.5)		(26.5)
慢性腎盂腎炎	344	474	818		818
(%)	(0.6)	(1.2)	(0.8)		(0.8)
急速進行性糸球体腎炎	433	512	945		945
(%)	(0.7)	(1.3)	(0.9)		(0.9)
妊娠腎/妊娠中毒症	2	271	273		273
(%)	(0.0)	(0.7)	(0.3)		(0.3)
その他の分類不能の腎炎	180	167	347		347
(%)	(0.3)	(0.4)	(0.3)		(0.3)
多発性嚢胞腎	1,156	1,203	2,359		2,359
(%)	(1.9)	(2.9)	(2.4)		(2.4)
腎硬化症	10,283	6,363	16,646		16,646
(%)	(17.3)	(15.5)	(16.6)		(16.6)
悪性高血圧	415	315	730		730
(%)	(0.7)	(0.8)	(0.7)		(0.7)
糖尿病性腎症	21,861	12,544	34,405		34,405
(%)	(36.8)	(30.6)	(34.3)		(34.3)
SLE腎炎	170	227	397		397
(%)	(0.3)	(0.6)	(0.4)		(0.4)
アミロイド腎	55	95	150		150
(%)	(0.1)	(0.2)	(0.1)		(0.1)
痛風腎	268	50	318		318
(%)	(0.5)	(0.1)	(0.3)		(0.3)
先天性代謝異常による腎不全	9	12	21		21
(%)	(0.0)	(0.0)	(0.0)		(0.0)
腎・尿路結核	40	36	76		76
(%)	(0.1)	(0.1)	(0.1)		(0.1)
腎・尿路結石	136	95	231		231
(%)	(0.2)	(0.2)	(0.2)		(0.2)
腎・尿路腫瘍	312	101	413		413
(%)	(0.5)	(0.2)	(0.4)		(0.4)
閉塞性尿路障害	148	75	223		223
(%)	(0.2)	(0.2)	(0.2)		(0.2)
骨髄腫	73	40	113		113
(%)	(0.1)	(0.1)	(0.1)		(0.1)
腎形成不全	27	34	61		61
(%)	(0.0)	(0.1)	(0.1)		(0.1)
不明	7,047	5,301	12,348		12,348
(%)	(11.9)	(12.9)	(12.3)		(12.3)
再導入	95	64	159		159
(%)	(0.2)	(0.2)	(0.2)		(0.2)
その他	1,508	1,170	2,678		2,678
(%)	(2.5)	(2.9)	(2.7)		(2.7)
合計	59,342	40,958	100,300		100,300
(%)	(100.0)	(100.0)	(100.0)		(100.0)
記載なし	4	4	8		8
総計	59,346	40,962	100,308		100,308

数値下のカッコ内は列方向の合計に対する%です。
 (C)Japanese Society for Dialysis Therapy
 2015年調査