

表 505 治療方法(新分類) 併用の状況別 (透析患者全体 75歳以上)

併用の状況	施設血液透析	Off-line HDF	On-line HDF	Push/Pull HDF	ハイオフィльтраーション	IHDF	血液濾過	血液吸着透析 (リクセル等使用)	在宅血液透析	腹膜透析 (CAPD)	腹膜透析 (APD)	腹膜透析 (CCPD)	合計	記載なし	総計
HD(F)単独	84,962	1,423	10,464	15	86	1,275	7	235	21				98,488		98,488
(%)	(100.0)	(99.8)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)				(98.2)		(98.2)
PD単独										1,072	471	78	1,621		1,621
(%)										(92.0)	(92.5)	(76.5)	(1.6)		(1.6)
HD(F)単独.PDカテあり	23	3	2										28		28
(%)	(0.0)	(0.2)	(0.0)										(0.0)		(0.0)
PD.HD併用週1										80	34	22	136		136
HD(F)										(6.9)	(6.7)	(21.6)	(0.1)		(0.1)
(%)															
PD.HD併用週2	2									5	2	1	10		10
HD(F)										(0.4)	(0.4)	(1.0)	(0.0)		(0.0)
(%)	(0.0)														
PD.HD併用週3	3												3		3
HD(F)													(0.0)		(0.0)
(%)	(0.0)														
PD.HD併用週4															
HD(F)															
(%)															
PD.HD併用その他	1		1							8	2	1	13		13
(%)	(0.0)		(0.0)							(0.7)	(0.4)	(1.0)	(0.0)		(0.0)
合計	84,991	1,426	10,467	15	86	1,275	7	235	21	1,165	509	102	100,299		100,299
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)
不明	7		1										8		8
記載なし	1												1		1
総計	84,999	1,426	10,468	15	86	1,275	7	235	21	1,165	509	102	100,308		100,308

数値下のかっこ内は列方向の合計に対する%です。
*腹膜透析 (CCPD) は、APDと手動のバック交換の両方を行うもの
(C)Japanese Society for Dialysis Therapy
2015年調査