

表 506 透析前補正カルシウム濃度(mg/dL) 透析前リン濃度(mg/dL)別 (透析患者全体 75歳以上)

透析前リン濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
3未満	34	14	79	228	81	972	1,557	1,247	189	510	276	134	64	50	5,435	88	5,523	9.43	0.88
(%)	(0.6)	(0.3)	(1.5)	(4.2)	(1.5)	(17.9)	(28.6)	(22.9)	(3.5)	(9.4)	(5.1)	(2.5)	(1.2)	(0.9)	(100.0)				
3.0～	14	20	67	343	111	1,442	1,915	1,237	155	418	208	72	32	24	6,058	65	6,123	9.27	0.73
(%)	(0.2)	(0.3)	(1.1)	(5.7)	(1.8)	(23.8)	(31.6)	(20.4)	(2.6)	(6.9)	(3.4)	(1.2)	(0.5)	(0.4)	(100.0)				
3.5～	29	41	166	592	173	2,298	3,067	1,898	245	533	261	96	41	33	9,473	111	9,584	9.22	0.73
(%)	(0.3)	(0.4)	(1.8)	(6.2)	(1.8)	(24.3)	(32.4)	(20.0)	(2.6)	(5.6)	(2.8)	(1.0)	(0.4)	(0.3)	(100.0)				
4.0～	38	49	197	824	240	3,359	4,201	2,510	293	669	341	133	51	36	12,941	176	13,117	9.20	0.71
(%)	(0.3)	(0.4)	(1.5)	(6.4)	(1.9)	(26.0)	(32.5)	(19.4)	(2.3)	(5.2)	(2.6)	(1.0)	(0.4)	(0.3)	(100.0)				
4.5～	39	52	237	947	263	3,749	4,580	2,575	300	797	361	131	46	36	14,113	161	14,274	9.18	0.70
(%)	(0.3)	(0.4)	(1.7)	(6.7)	(1.9)	(26.6)	(32.5)	(18.2)	(2.1)	(5.6)	(2.6)	(0.9)	(0.3)	(0.3)	(100.0)				
5.0～	39	56	254	968	269	3,369	4,124	2,536	291	735	364	103	56	30	13,194	128	13,322	9.18	0.71
(%)	(0.3)	(0.4)	(1.9)	(7.3)	(2.0)	(25.5)	(31.3)	(19.2)	(2.2)	(5.6)	(2.8)	(0.8)	(0.4)	(0.2)	(100.0)				
5.5～	38	51	241	858	244	2,669	3,210	2,019	227	603	292	106	28	24	10,610	118	10,728	9.16	0.73
(%)	(0.4)	(0.5)	(2.3)	(8.1)	(2.3)	(25.2)	(30.3)	(19.0)	(2.1)	(5.7)	(2.8)	(1.0)	(0.3)	(0.2)	(100.0)				
6.0～	6	5	41	132	46	443	529	358	37	88	53	21	5	3	1,767	14	1,781	9.18	0.72
(%)	(0.3)	(0.3)	(2.3)	(7.5)	(2.6)	(25.1)	(29.9)	(20.3)	(2.1)	(5.0)	(3.0)	(1.2)	(0.3)	(0.2)	(100.0)				
6.1～	25	28	133	471	127	1,307	1,584	1,055	139	341	178	59	15	15	5,477	96	5,573	9.18	0.75
(%)	(0.5)	(0.5)	(2.4)	(8.6)	(2.3)	(23.9)	(28.9)	(19.3)	(2.5)	(6.2)	(3.2)	(1.1)	(0.3)	(0.3)	(100.0)				
6.5～	27	23	137	410	109	1,119	1,188	778	127	269	142	49	12	12	4,402	52	4,454	9.14	0.78
(%)	(0.6)	(0.5)	(3.1)	(9.3)	(2.5)	(25.4)	(27.0)	(17.7)	(2.9)	(6.1)	(3.2)	(1.1)	(0.3)	(0.3)	(100.0)				
7.0～	14	20	86	238	64	607	687	482	63	157	77	27	14	4	2,540	36	2,576	9.15	0.78
(%)	(0.6)	(0.8)	(3.4)	(9.4)	(2.5)	(23.9)	(27.0)	(19.0)	(2.5)	(6.2)	(3.0)	(1.1)	(0.6)	(0.2)	(100.0)				
7.5～	6	7	61	155	23	356	395	241	25	71	53	21	11	4	1,429	25	1,454	9.13	0.80
(%)	(0.4)	(0.5)	(4.3)	(10.8)	(1.6)	(24.9)	(27.6)	(16.9)	(1.7)	(5.0)	(3.7)	(1.5)	(0.8)	(0.3)	(100.0)				
8.0～	6	8	24	71	23	173	207	125	21	56	24	10	7	1	756	18	774	9.15	0.86
(%)	(0.8)	(1.1)	(3.2)	(9.4)	(3.0)	(22.9)	(27.4)	(16.5)	(2.8)	(7.4)	(3.2)	(1.3)	(0.9)	(0.1)	(100.0)				
8.5～	4	4	13	42	9	121	87	75	7	24	14	6	4	1	411	4	415	9.10	0.88
(%)	(1.0)	(1.0)	(3.2)	(10.2)	(2.2)	(29.4)	(21.2)	(18.2)	(1.7)	(5.8)	(3.4)	(1.5)	(1.0)	(0.2)	(100.0)				
9.0～	9	9	29	58	15	135	149	96	16	32	26	7	2	2	585	14	599	9.07	0.92
(%)	(1.5)	(1.5)	(5.0)	(9.9)	(2.6)	(23.1)	(25.5)	(16.4)	(2.7)	(5.5)	(4.4)	(1.2)	(0.3)	(0.3)	(100.0)				
合計	328	387	1,765	6,337	1,797	22,119	27,480	17,232	2,135	5,303	2,670	975	388	275	89,191	1,106	90,297	9.20	0.74
(%)	(0.4)	(0.4)	(2.0)	(7.1)	(2.0)	(24.8)	(30.8)	(19.3)	(2.4)	(5.9)	(3.0)	(1.1)	(0.4)	(0.3)	(100.0)				
記載なし	2		2	7	2	23	46	43	6	9	9	5	4	3	161	9,850	10,011	9.50	0.96
(%)	(1.2)		(1.2)	(4.3)	(1.2)	(14.3)	(28.6)	(26.7)	(3.7)	(5.6)	(5.6)	(3.1)	(2.5)	(1.9)	(100.0)				
総計	330	387	1,767	6,344	1,799	22,142	27,526	17,275	2,141	5,312	2,679	980	392	278	89,352	10,956	100,308	9.20	0.74
(%)	(0.4)	(0.4)	(2.0)	(7.1)	(2.0)	(24.8)	(30.8)	(19.3)	(2.4)	(5.9)	(3.0)	(1.1)	(0.4)	(0.3)	(100.0)				
平均	5.14	5.24	5.30	5.15	5.05	4.94	4.84	4.83	4.81	4.80	4.82	4.70	4.54	4.34	4.89	4.93	4.89		
標準偏差	1.74	1.54	1.49	1.36	1.35	1.28	1.29	1.35	1.42	1.43	1.55	1.55	1.70	1.65	1.34	1.49	1.35		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査