

表 509 透析前補正カルシウム 透析前リン濃度(透析患者全体、糖尿病あり患者)

透析前リン濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
3未満	44	18	67	245	71	1,005	1,560	1,180	188	485	281	130	59	44	5,377	80	5,457	9.41	0.88
(%)	(0.8)	(0.3)	(1.2)	(4.6)	(1.3)	(18.7)	(29.0)	(21.9)	(3.5)	(9.0)	(5.2)	(2.4)	(1.1)	(0.8)	(100.0)				
3.0～	19	13	99	397	117	1,549	2,026	1,238	169	392	211	76	29	26	6,361	59	6,420	9.24	0.74
(%)	(0.3)	(0.2)	(1.6)	(6.2)	(1.8)	(24.4)	(31.9)	(19.5)	(2.7)	(6.2)	(3.3)	(1.2)	(0.5)	(0.4)	(100.0)				
3.5～	39	46	196	712	206	2,800	3,620	2,158	224	578	296	85	39	31	11,030	120	11,150	9.19	0.71
(%)	(0.4)	(0.4)	(1.8)	(6.5)	(1.9)	(25.4)	(32.8)	(19.6)	(2.0)	(5.2)	(2.7)	(0.8)	(0.4)	(0.3)	(100.0)				
4.0～	43	53	281	1,167	359	4,278	5,230	3,054	330	774	393	121	39	36	16,158	192	16,350	9.16	0.69
(%)	(0.3)	(0.3)	(1.7)	(7.2)	(2.2)	(26.5)	(32.4)	(18.9)	(2.0)	(4.8)	(2.4)	(0.7)	(0.2)	(0.2)	(100.0)				
4.5～	67	90	440	1,481	415	5,217	6,076	3,310	385	956	445	124	48	39	19,093	209	19,302	9.13	0.70
(%)	(0.4)	(0.5)	(2.3)	(7.8)	(2.2)	(27.3)	(31.8)	(17.3)	(2.0)	(5.0)	(2.3)	(0.6)	(0.3)	(0.2)	(100.0)				
5.0～	60	109	469	1,621	474	5,330	6,037	3,467	354	969	452	134	46	29	19,551	189	19,740	9.12	0.70
(%)	(0.3)	(0.6)	(2.4)	(8.3)	(2.4)	(27.3)	(30.9)	(17.7)	(1.8)	(5.0)	(2.3)	(0.7)	(0.2)	(0.1)	(100.0)				
5.5～	57	116	455	1,450	398	4,530	4,974	2,930	318	834	384	128	44	27	16,645	180	16,825	9.11	0.71
(%)	(0.3)	(0.7)	(2.7)	(8.7)	(2.4)	(27.2)	(29.9)	(17.6)	(1.9)	(5.0)	(2.3)	(0.8)	(0.3)	(0.2)	(100.0)				
6.0～	16	21	78	262	72	720	945	520	49	142	70	19	3	6	2,923	27	2,950	9.10	0.73
(%)	(0.5)	(0.7)	(2.7)	(9.0)	(2.5)	(24.6)	(32.3)	(17.8)	(1.7)	(4.9)	(2.4)	(0.7)	(0.1)	(0.2)	(100.0)				
6.1～	51	87	325	976	261	2,633	2,891	1,761	220	491	259	71	22	12	10,060	143	10,203	9.09	0.73
(%)	(0.5)	(0.9)	(3.2)	(9.7)	(2.6)	(26.2)	(28.7)	(17.5)	(2.2)	(4.9)	(2.6)	(0.7)	(0.2)	(0.1)	(100.0)				
6.5～	48	81	310	865	260	2,351	2,501	1,511	207	427	200	68	17	3	8,849	109	8,958	9.06	0.73
(%)	(0.5)	(0.9)	(3.5)	(9.8)	(2.9)	(26.6)	(28.3)	(17.1)	(2.3)	(4.8)	(2.3)	(0.8)	(0.2)	(0.0)	(100.0)				
7.0～	45	62	250	616	146	1,435	1,520	903	117	266	126	49	12	6	5,553	82	5,635	9.03	0.77
(%)	(0.8)	(1.1)	(4.5)	(11.1)	(2.6)	(25.8)	(27.4)	(16.3)	(2.1)	(4.8)	(2.3)	(0.9)	(0.2)	(0.1)	(100.0)				
7.5～	26	55	167	369	92	883	940	562	77	169	98	24	6	9	3,477	57	3,534	9.03	0.79
(%)	(0.7)	(1.6)	(4.8)	(10.6)	(2.6)	(25.4)	(27.0)	(16.2)	(2.2)	(4.9)	(2.8)	(0.7)	(0.2)	(0.3)	(100.0)				
8.0～	21	31	91	251	76	520	544	340	36	105	40	19	8	4	2,086	37	2,123	9.00	0.81
(%)	(1.0)	(1.5)	(4.4)	(12.0)	(3.6)	(24.9)	(26.1)	(16.3)	(1.7)	(5.0)	(1.9)	(0.9)	(0.4)	(0.2)	(100.0)				
8.5～	22	23	70	175	29	365	314	207	23	53	25	11	2	2	1,321	25	1,346	8.93	0.83
(%)	(1.7)	(1.7)	(5.3)	(13.2)	(2.2)	(27.6)	(23.8)	(15.7)	(1.7)	(4.0)	(1.9)	(0.8)	(0.2)	(0.2)	(100.0)				
9.0～	41	57	135	272	58	450	487	280	34	88	46	11	7	3	1,969	50	2,019	8.88	0.90
(%)	(2.1)	(2.9)	(6.9)	(13.8)	(2.9)	(22.9)	(24.7)	(14.2)	(1.7)	(4.5)	(2.3)	(0.6)	(0.4)	(0.2)	(100.0)				
合計	599	862	3,433	10,859	3,034	34,066	39,665	23,421	2,731	6,729	3,326	1,070	381	277	130,453	1,559	132,012	9.13	0.73
(%)	(0.5)	(0.7)	(2.6)	(8.3)	(2.3)	(26.1)	(30.4)	(18.0)	(2.1)	(5.2)	(2.5)	(0.8)	(0.3)	(0.2)	(100.0)				
記載なし	6	2	4	13	2	34	59	49	12	16	15	3	2	3	220	12,638	12,858	9.31	1.07
(%)	(2.7)	(0.9)	(1.8)	(5.9)	(0.9)	(15.5)	(26.8)	(22.3)	(5.5)	(7.3)	(6.8)	(1.4)	(0.9)	(1.4)	(100.0)				
総計	605	864	3,437	10,872	3,036	34,100	39,724	23,470	2,743	6,745	3,341	1,073	383	280	130,673	14,197	144,870	9.13	0.73
(%)	(0.5)	(0.7)	(2.6)	(8.3)	(2.3)	(26.1)	(30.4)	(18.0)	(2.1)	(5.2)	(2.6)	(0.8)	(0.3)	(0.2)	(100.0)				
平均	5.83	6.10	5.78	5.53	5.45	5.26	5.16	5.14	5.12	5.09	5.03	4.94	4.69	4.49	5.23	5.41	5.24		
標準偏差	2.04	1.80	1.60	1.50	1.44	1.39	1.40	1.44	1.52	1.50	1.57	1.66	1.74	1.69	1.45	1.68	1.46		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査