

表 549 HDF希釈方法 透析後体重(kg)別（オンラインHDF 女性のみ 透析回数週3回 透析歴2年以上）

透析後体重	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
20未満	3				3			3
(%)	(100.0)				(100.0)			
20～	49	2			51		6	57
(%)	(96.1)	(3.9)			(100.0)			
30～	1,743	96		10	1,849		158	2,007
(%)	(94.3)	(5.2)		(0.5)	(100.0)			
40～	4,829	249		28	5,106		423	5,529
(%)	(94.6)	(4.9)		(0.5)	(100.0)			
50～	2,920	142		17	3,079		257	3,336
(%)	(94.8)	(4.6)		(0.6)	(100.0)			
60～	1,029	40		4	1,073		98	1,171
(%)	(95.9)	(3.7)		(0.4)	(100.0)			
70～	345	19			364		33	397
(%)	(94.8)	(5.2)			(100.0)			
80～	121	6		1	128		11	139
(%)	(94.5)	(4.7)		(0.8)	(100.0)			
90～	29	1		1	31		1	32
(%)	(93.5)	(3.2)		(3.2)	(100.0)			
100～	11				11			11
(%)	(100.0)				(100.0)			
110～	3				3			3
(%)	(100.0)				(100.0)			
120～	1				1			1
(%)	(100.0)				(100.0)			
合計	11,083	555		61	11,699		987	12,686
(%)	(94.7)	(4.7)		(0.5)	(100.0)			
記載なし	195	16		37	248		53	301
(%)	(78.6)	(6.5)		(14.9)	(100.0)			
総計	11,278	571		98	11,947		1,040	12,987
(%)	(94.4)	(4.8)		(0.8)	(100.0)			
平均	49.35	48.67		48.99	49.31		49.27	49.31
標準偏差	10.53	10.36		10.93	10.52		10.29	10.50

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査