

表 555 糖尿病の既往 治療方法(新分類)別 (透析患者全体)

治療方法 (新分類)	既往なし	既往あり	合計	不明	記載なし	総計
施設血液透析	98,035	115,680	213,715	177	34,833	248,725
(%)	(45.9)	(54.1)	(100.0)			
Off-line HDF	2,399	1,974	4,373	2	957	5,332
(%)	(54.9)	(45.1)	(100.0)			
On-line HDF	19,614	18,125	37,739	17	6,771	44,527
(%)	(52.0)	(48.0)	(100.0)			
Push/Pull HDF	43	48	91		19	110
(%)	(47.3)	(52.7)	(100.0)			
ハイオフィльтраーション	112	123	235		32	267
(%)	(47.7)	(52.3)	(100.0)			
IHDF	1,424	1,704	3,128	1	411	3,540
(%)	(45.5)	(54.5)	(100.0)			
血液濾過	6	6	12		4	16
(%)	(50.0)	(50.0)	(100.0)			
血液吸着透析 (リクセル等使用)	1,022	138	1,160	1	215	1,376
(%)	(88.1)	(11.9)	(100.0)			
在宅血液透析	337	116	453		102	555
(%)	(74.4)	(25.6)	(100.0)			
腹膜透析 (CAPD)	1,950	1,842	3,792	7	1,103	4,902
(%)	(51.4)	(48.6)	(100.0)			
腹膜透析 (APD)	1,093	1,109	2,202	6	567	2,775
(%)	(49.6)	(50.4)	(100.0)			
腹膜透析 (CCPD)	455	392	847	4	241	1,092
(%)	(53.7)	(46.3)	(100.0)			
合計	126,490	141,257	267,747	215	45,255	313,217
(%)	(47.2)	(52.8)	(100.0)			
記載なし						
(%)						
総計	126,490	141,257	267,747	215	45,255	313,217
(%)	(47.2)	(52.8)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査