

表 560 糖尿病の既往 原疾患別（透析患者全体）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	63,033	9,616	72,649	59	20,639	93,347
(%)	(86.8)	(13.2)	(100.0)			
慢性腎盂腎炎	2,002	338	2,340	2	593	2,935
(%)	(85.6)	(14.4)	(100.0)			
急速進行性糸球体腎炎	1,494	480	1,974	3	501	2,478
(%)	(75.7)	(24.3)	(100.0)			
妊娠腎/妊娠中毒症	1,103	115	1,218	1	326	1,545
(%)	(90.6)	(9.4)	(100.0)			
その他の分類不能の腎炎	924	163	1,087	2	316	1,405
(%)	(85.0)	(15.0)	(100.0)			
多発性嚢胞腎	8,047	834	8,881	6	2,369	11,256
(%)	(90.6)	(9.4)	(100.0)			
腎硬化症	19,329	4,410	23,739	39	6,027	29,805
(%)	(81.4)	(18.6)	(100.0)			
悪性高血圧	1,754	299	2,053	5	560	2,618
(%)	(85.4)	(14.6)	(100.0)			
糖尿病性腎症	12	116,665	116,677		3,601	120,278
(%)	(0.0)	(100.0)	(100.0)			
SLE腎炎	1,476	304	1,780	1	446	2,227
(%)	(82.9)	(17.1)	(100.0)			
アミロイド腎	278	78	356		106	462
(%)	(78.1)	(21.9)	(100.0)			
痛風腎	747	106	853		222	1,075
(%)	(87.6)	(12.4)	(100.0)			
先天性代謝異常による腎不全	171	32	203		71	274
(%)	(84.2)	(15.8)	(100.0)			
腎・尿路結核	131	19	150		45	195
(%)	(87.3)	(12.7)	(100.0)			
腎・尿路結石	407	60	467		114	581
(%)	(87.2)	(12.8)	(100.0)			
腎・尿路腫瘍	591	117	708	1	201	910
(%)	(83.5)	(16.5)	(100.0)			
閉塞性尿路障害	519	65	584		146	730
(%)	(88.9)	(11.1)	(100.0)			
骨髄腫	177	43	220	2	69	291
(%)	(80.5)	(19.5)	(100.0)			
腎形成不全	486	40	526	1	133	660
(%)	(92.4)	(7.6)	(100.0)			
不明	17,836	5,417	23,253	79	6,565	29,897
(%)	(76.7)	(23.3)	(100.0)			
再導入	1,434	311	1,745	1	437	2,183
(%)	(82.2)	(17.8)	(100.0)			
その他	4,528	1,742	6,270	13	1,761	8,044
(%)	(72.2)	(27.8)	(100.0)			
合計	126,479	141,254	267,733	215	45,248	313,196
(%)	(47.2)	(52.8)	(100.0)			
記載なし	11	3	14		7	21
(%)	(78.6)	(21.4)	(100.0)			
総計	126,490	141,257	267,747	215	45,255	313,217
(%)	(47.2)	(52.8)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査