

表 591 糖尿病の既往 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	45,640	7,087	52,727	45	10,702	63,474
(%)	(86.6)	(13.4)	(100.0)			
慢性腎盂腎炎	1,468	255	1,723	2	315	2,040
(%)	(85.2)	(14.8)	(100.0)			
急速進行性糸球体腎炎	1,142	374	1,516	3	252	1,771
(%)	(75.3)	(24.7)	(100.0)			
妊娠腎/妊娠中毒症	738	84	822		159	981
(%)	(89.8)	(10.2)	(100.0)			
その他の分類不能の腎炎	641	127	768	2	159	929
(%)	(83.5)	(16.5)	(100.0)			
多発性嚢胞腎	6,148	600	6,748	4	1,264	8,016
(%)	(91.1)	(8.9)	(100.0)			
腎硬化症	15,289	3,386	18,675	30	3,302	22,007
(%)	(81.9)	(18.1)	(100.0)			
悪性高血圧	1,341	217	1,558	5	298	1,861
(%)	(86.1)	(13.9)	(100.0)			
糖尿病性腎症	10	87,892	87,902		1,524	89,426
(%)	(0.0)	(100.0)	(100.0)			
SLE腎炎	1,056	231	1,287		217	1,504
(%)	(82.1)	(17.9)	(100.0)			
アミロイド腎	197	53	250		64	314
(%)	(78.8)	(21.2)	(100.0)			
痛風腎	575	87	662		116	778
(%)	(86.9)	(13.1)	(100.0)			
先天性代謝異常による腎不全	121	25	146		27	173
(%)	(82.9)	(17.1)	(100.0)			
腎・尿路結核	102	15	117		24	141
(%)	(87.2)	(12.8)	(100.0)			
腎・尿路結石	315	47	362		59	421
(%)	(87.0)	(13.0)	(100.0)			
腎・尿路腫瘍	483	86	569		109	678
(%)	(84.9)	(15.1)	(100.0)			
閉塞性尿路障害	376	48	424		70	494
(%)	(88.7)	(11.3)	(100.0)			
骨髄腫	126	27	153	1	36	190
(%)	(82.4)	(17.6)	(100.0)			
腎形成不全	317	26	343	1	56	400
(%)	(92.4)	(7.6)	(100.0)			
不明	13,542	4,025	17,567	51	3,558	21,176
(%)	(77.1)	(22.9)	(100.0)			
再導入	950	234	1,184	1	171	1,356
(%)	(80.2)	(19.8)	(100.0)			
その他	3,414	1,338	4,752	7	956	5,715
(%)	(71.8)	(28.2)	(100.0)			
合計	93,991	106,264	200,255	152	23,438	223,845
(%)	(46.9)	(53.1)	(100.0)			
記載なし	6	1	7			7
(%)	(85.7)	(14.3)	(100.0)			
総計	93,997	106,265	200,262	152	23,438	223,852
(%)	(46.9)	(53.1)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査