

表 665 糖尿病の既往 原疾患別（腹膜透析患者）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	1,730	266	1,996	5	829	2,830
(%)	(86.7)	(13.3)	(100.0)			
慢性腎盂腎炎	68	9	77		20	97
(%)	(88.3)	(11.7)	(100.0)			
急速進行性糸球体腎炎	32	11	43		23	66
(%)	(74.4)	(25.6)	(100.0)			
妊娠腎/妊娠中毒症	21	3	24		3	27
(%)	(87.5)	(12.5)	(100.0)			
その他の分類不能の腎炎	50	6	56		22	78
(%)	(89.3)	(10.7)	(100.0)			
多発性嚢胞腎	157	22	179		84	263
(%)	(87.7)	(12.3)	(100.0)			
腎硬化症	598	163	761	2	332	1,095
(%)	(78.6)	(21.4)	(100.0)			
悪性高血圧	61	14	75		32	107
(%)	(81.3)	(18.7)	(100.0)			
糖尿病性腎症	1	2,638	2,639		160	2,799
(%)	(0.0)	(100.0)	(100.0)			
SLE腎炎	28	5	33	1	12	46
(%)	(84.8)	(15.2)	(100.0)			
アミロイド腎	9	1	10		5	15
(%)	(90.0)	(10.0)	(100.0)			
痛風腎	22	4	26		6	32
(%)	(84.6)	(15.4)	(100.0)			
先天性代謝異常による腎不全	10	1	11		6	17
(%)	(90.9)	(9.1)	(100.0)			
腎・尿路結核					2	2
(%)						
腎・尿路結石	15		15		6	21
(%)	(100.0)		(100.0)			
腎・尿路腫瘍	8	3	11	1	2	14
(%)	(72.7)	(27.3)	(100.0)			
閉塞性尿路障害	13	1	14		7	21
(%)	(92.9)	(7.1)	(100.0)			
骨髄腫	2	1	3		4	7
(%)	(66.7)	(33.3)	(100.0)			
腎形成不全	47	5	52		20	72
(%)	(90.4)	(9.6)	(100.0)			
不明	459	137	596	7	239	842
(%)	(77.0)	(23.0)	(100.0)			
再導入	32	7	39		16	55
(%)	(82.1)	(17.9)	(100.0)			
その他	135	46	181	1	81	263
(%)	(74.6)	(25.4)	(100.0)			
合計	3,498	3,343	6,841	17	1,911	8,769
(%)	(51.1)	(48.9)	(100.0)			
記載なし						
(%)						
総計	3,498	3,343	6,841	17	1,911	8,769
(%)	(51.1)	(48.9)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査