

表 693 心筋梗塞の既往 治療方法(新分類)別 (透析患者全体)

治療方法 (新分類)	既往なし	既往あり	合計	不明	記載なし	総計
施設血液透析	169,464	19,586	189,050	706	58,969	248,725
(%)	(89.6)	(10.4)	(100.0)			
Off-line HDF	3,370	618	3,988	16	1,328	5,332
(%)	(84.5)	(15.5)	(100.0)			
On-line HDF	30,174	3,350	33,524	39	10,964	44,527
(%)	(90.0)	(10.0)	(100.0)			
Push/Pull HDF	72	10	82		28	110
(%)	(87.8)	(12.2)	(100.0)			
ハイオフィльтраーション	170	45	215	1	51	267
(%)	(79.1)	(20.9)	(100.0)			
IHDF	2,473	310	2,783	13	744	3,540
(%)	(88.9)	(11.1)	(100.0)			
血液濾過	9	2	11		5	16
(%)	(81.8)	(18.2)	(100.0)			
血液吸着透析 (リクセル等使用)	1,035	108	1,143		233	1,376
(%)	(90.6)	(9.4)	(100.0)			
在宅血液透析	380	21	401	1	153	555
(%)	(94.8)	(5.2)	(100.0)			
腹膜透析 (CAPD)	2,880	268	3,148	10	1,744	4,902
(%)	(91.5)	(8.5)	(100.0)			
腹膜透析 (APD)	1,742	156	1,898	10	867	2,775
(%)	(91.8)	(8.2)	(100.0)			
腹膜透析 (CCPD)	695	64	759	1	332	1,092
(%)	(91.6)	(8.4)	(100.0)			
合計	212,464	24,538	237,002	797	75,418	313,217
(%)	(89.6)	(10.4)	(100.0)			
記載なし						
(%)						
総計	212,464	24,538	237,002	797	75,418	313,217
(%)	(89.6)	(10.4)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析 (CCPD) は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査