

表 698 心筋梗塞の既往 原疾患別（透析患者全体）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	65,135	5,322	70,457	209	22,681	93,347
(%)	(92.4)	(7.6)	(100.0)			
慢性腎盂腎炎	2,104	152	2,256	5	674	2,935
(%)	(93.3)	(6.7)	(100.0)			
急速進行性糸球体腎炎	1,770	121	1,891	8	579	2,478
(%)	(93.6)	(6.4)	(100.0)			
妊娠腎/妊娠中毒症	1,116	67	1,183	4	358	1,545
(%)	(94.3)	(5.7)	(100.0)			
その他の分類不能の腎炎	964	70	1,034	5	366	1,405
(%)	(93.2)	(6.8)	(100.0)			
多発性嚢胞腎	8,124	535	8,659	23	2,574	11,256
(%)	(93.8)	(6.2)	(100.0)			
腎硬化症	20,611	2,362	22,973	45	6,787	29,805
(%)	(89.7)	(10.3)	(100.0)			
悪性高血圧	1,833	162	1,995	6	617	2,618
(%)	(91.9)	(8.1)	(100.0)			
糖尿病性腎症	78,525	12,946	91,471	313	28,494	120,278
(%)	(85.8)	(14.2)	(100.0)			
SLE腎炎	1,614	102	1,716	7	504	2,227
(%)	(94.1)	(5.9)	(100.0)			
アミロイド腎	313	20	333		129	462
(%)	(94.0)	(6.0)	(100.0)			
痛風腎	773	56	829	2	244	1,075
(%)	(93.2)	(6.8)	(100.0)			
先天性代謝異常による腎不全	181	8	189	1	84	274
(%)	(95.8)	(4.2)	(100.0)			
腎・尿路結核	140	9	149		46	195
(%)	(94.0)	(6.0)	(100.0)			
腎・尿路結石	425	26	451		130	581
(%)	(94.2)	(5.8)	(100.0)			
腎・尿路腫瘍	646	41	687	1	222	910
(%)	(94.0)	(6.0)	(100.0)			
閉塞性尿路障害	534	30	564	2	164	730
(%)	(94.7)	(5.3)	(100.0)			
骨髄腫	196	15	211	1	79	291
(%)	(92.9)	(7.1)	(100.0)			
腎形成不全	493	17	510		150	660
(%)	(96.7)	(3.3)	(100.0)			
不明	20,053	1,848	21,901	125	7,871	29,897
(%)	(91.6)	(8.4)	(100.0)			
再導入	1,518	123	1,641	3	539	2,183
(%)	(92.5)	(7.5)	(100.0)			
その他	5,383	505	5,888	37	2,119	8,044
(%)	(91.4)	(8.6)	(100.0)			
合計	212,451	24,537	236,988	797	75,411	313,196
(%)	(89.6)	(10.4)	(100.0)			
記載なし	13	1	14		7	21
(%)	(92.9)	(7.1)	(100.0)			
総計	212,464	24,538	237,002	797	75,418	313,217
(%)	(89.6)	(10.4)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査