

表 728 心筋梗塞の既往 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	47,515	3,827	51,342	178	11,954	63,474
(%)	(92.5)	(7.5)	(100.0)			
慢性腎盂腎炎	1,556	106	1,662	4	374	2,040
(%)	(93.6)	(6.4)	(100.0)			
急速進行性糸球体腎炎	1,372	91	1,463	7	301	1,771
(%)	(93.8)	(6.2)	(100.0)			
妊娠腎/妊娠中毒症	757	47	804	4	173	981
(%)	(94.2)	(5.8)	(100.0)			
その他の分類不能の腎炎	680	53	733	5	191	929
(%)	(92.8)	(7.2)	(100.0)			
多発性嚢胞腎	6,208	384	6,592	19	1,405	8,016
(%)	(94.2)	(5.8)	(100.0)			
腎硬化症	16,420	1,806	18,226	35	3,746	22,007
(%)	(90.1)	(9.9)	(100.0)			
悪性高血圧	1,425	105	1,530	4	327	1,861
(%)	(93.1)	(6.9)	(100.0)			
糖尿病性腎症	62,924	10,056	72,980	271	16,175	89,426
(%)	(86.2)	(13.8)	(100.0)			
SLE腎炎	1,167	73	1,240	6	258	1,504
(%)	(94.1)	(5.9)	(100.0)			
アミロイド腎	221	15	236		78	314
(%)	(93.6)	(6.4)	(100.0)			
痛風腎	604	38	642	2	134	778
(%)	(94.1)	(5.9)	(100.0)			
先天性代謝異常による腎不全	131	6	137	1	35	173
(%)	(95.6)	(4.4)	(100.0)			
腎・尿路結核	107	8	115		26	141
(%)	(93.0)	(7.0)	(100.0)			
腎・尿路結石	334	16	350		71	421
(%)	(95.4)	(4.6)	(100.0)			
腎・尿路腫瘍	521	33	554	1	123	678
(%)	(94.0)	(6.0)	(100.0)			
閉塞性尿路障害	393	18	411	2	81	494
(%)	(95.6)	(4.4)	(100.0)			
骨髄腫	137	13	150		40	190
(%)	(91.3)	(8.7)	(100.0)			
腎形成不全	321	13	334		66	400
(%)	(96.1)	(3.9)	(100.0)			
不明	15,352	1,415	16,767	102	4,307	21,176
(%)	(91.6)	(8.4)	(100.0)			
再導入	1,044	78	1,122	2	232	1,356
(%)	(93.0)	(7.0)	(100.0)			
その他	4,100	384	4,484	33	1,198	5,715
(%)	(91.4)	(8.6)	(100.0)			
合計	163,289	18,585	181,874	676	41,295	223,845
(%)	(89.8)	(10.2)	(100.0)			
記載なし	7		7			7
(%)	(100.0)		(100.0)			
総計	163,296	18,585	181,881	676	41,295	223,852
(%)	(89.8)	(10.2)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査