

表 764 心筋梗塞の既往 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	13,046	1,085	14,131	19	3,312	17,462
(%)	(92.3)	(7.7)	(100.0)			
慢性腎盂腎炎	363	32	395	1	88	484
(%)	(91.9)	(8.1)	(100.0)			
急速進行性糸球体腎炎	274	21	295	1	57	353
(%)	(92.9)	(7.1)	(100.0)			
妊娠腎/妊娠中毒症	286	15	301		100	401
(%)	(95.0)	(5.0)	(100.0)			
その他の分類不能の腎炎	190	12	202		59	261
(%)	(94.1)	(5.9)	(100.0)			
多発性嚢胞腎	1,405	113	1,518	3	359	1,880
(%)	(92.6)	(7.4)	(100.0)			
腎硬化症	2,704	358	3,062	3	755	3,820
(%)	(88.3)	(11.7)	(100.0)			
悪性高血圧	280	40	320	1	70	391
(%)	(87.5)	(12.5)	(100.0)			
糖尿病性腎症	11,534	2,023	13,557	23	3,380	16,960
(%)	(85.1)	(14.9)	(100.0)			
SLE腎炎	340	20	360	1	65	426
(%)	(94.4)	(5.6)	(100.0)			
アミロイド腎	61	5	66		8	74
(%)	(92.4)	(7.6)	(100.0)			
痛風腎	122	11	133		33	166
(%)	(91.7)	(8.3)	(100.0)			
先天性代謝異常による腎不全	32	1	33		16	49
(%)	(97.0)	(3.0)	(100.0)			
腎・尿路結核	25		25		6	31
(%)	(100.0)		(100.0)			
腎・尿路結石	56	7	63		16	79
(%)	(88.9)	(11.1)	(100.0)			
腎・尿路腫瘍	94	5	99		20	119
(%)	(94.9)	(5.1)	(100.0)			
閉塞性尿路障害	76	10	86		18	104
(%)	(88.4)	(11.6)	(100.0)			
骨髄腫	24	1	25	1	7	33
(%)	(96.0)	(4.0)	(100.0)			
腎形成不全	109	3	112		24	136
(%)	(97.3)	(2.7)	(100.0)			
不明	3,261	297	3,558	7	1,138	4,703
(%)	(91.7)	(8.3)	(100.0)			
再導入	359	32	391	1	91	483
(%)	(91.8)	(8.2)	(100.0)			
その他	840	74	914	3	243	1,160
(%)	(91.9)	(8.1)	(100.0)			
合計	35,481	4,165	39,646	64	9,865	49,575
(%)	(89.5)	(10.5)	(100.0)			
記載なし	5	1	6			6
(%)	(83.3)	(16.7)	(100.0)			
総計	35,486	4,166	39,652	64	9,865	49,581
(%)	(89.5)	(10.5)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査