

表 800 心筋梗塞の既往 原疾患別（腹膜透析患者）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	1,800	90	1,890	5	935	2,830
(%)	(95.2)	(4.8)	(100.0)			
慢性腎盂腎炎	72	3	75		22	97
(%)	(96.0)	(4.0)	(100.0)			
急速進行性糸球体腎炎	37	2	39		27	66
(%)	(94.9)	(5.1)	(100.0)			
妊娠腎/妊娠中毒症	22	2	24		3	27
(%)	(91.7)	(8.3)	(100.0)			
その他の分類不能の腎炎	53	1	54		24	78
(%)	(98.1)	(1.9)	(100.0)			
多発性嚢胞腎	161	7	168		95	263
(%)	(95.8)	(4.2)	(100.0)			
腎硬化症	658	73	731	2	362	1,095
(%)	(90.0)	(10.0)	(100.0)			
悪性高血圧	62	5	67		40	107
(%)	(92.5)	(7.5)	(100.0)			
糖尿病性腎症	1,599	249	1,848	6	945	2,799
(%)	(86.5)	(13.5)	(100.0)			
SLE腎炎	30	1	31		15	46
(%)	(96.8)	(3.2)	(100.0)			
アミロイド腎	10		10		5	15
(%)	(100.0)		(100.0)			
痛風腎	25	1	26		6	32
(%)	(96.2)	(3.8)	(100.0)			
先天性代謝異常による腎不全	10		10		7	17
(%)	(100.0)		(100.0)			
腎・尿路結核					2	2
(%)						
腎・尿路結石	15	1	16		5	21
(%)	(93.8)	(6.3)	(100.0)			
腎・尿路腫瘍	10	1	11		3	14
(%)	(90.9)	(9.1)	(100.0)			
閉塞性尿路障害	12	2	14		7	21
(%)	(85.7)	(14.3)	(100.0)			
骨髄腫	2	1	3		4	7
(%)	(66.7)	(33.3)	(100.0)			
腎形成不全	48	1	49		23	72
(%)	(98.0)	(2.0)	(100.0)			
不明	514	32	546	7	289	842
(%)	(94.1)	(5.9)	(100.0)			
再導入	26	3	29		26	55
(%)	(89.7)	(10.3)	(100.0)			
その他	151	13	164	1	98	263
(%)	(92.1)	(7.9)	(100.0)			
合計	5,317	488	5,805	21	2,943	8,769
(%)	(91.6)	(8.4)	(100.0)			
記載なし						
(%)						
総計	5,317	488	5,805	21	2,943	8,769
(%)	(91.6)	(8.4)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査