

表 832 脳出血の既往 原疾患別（透析患者全体）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	65,805	4,534	70,339	186	22,822	93,347
(%)	(93.6)	(6.4)	(100.0)			
慢性腎盂腎炎	2,124	130	2,254	4	677	2,935
(%)	(94.2)	(5.8)	(100.0)			
急速進行性糸球体腎炎	1,804	88	1,892	9	577	2,478
(%)	(95.3)	(4.7)	(100.0)			
妊娠腎/妊娠中毒症	1,097	80	1,177	3	365	1,545
(%)	(93.2)	(6.8)	(100.0)			
その他の分類不能の腎炎	986	56	1,042	3	360	1,405
(%)	(94.6)	(5.4)	(100.0)			
多発性嚢胞腎	7,629	1,106	8,735	14	2,507	11,256
(%)	(87.3)	(12.7)	(100.0)			
腎硬化症	21,177	1,660	22,837	63	6,905	29,805
(%)	(92.7)	(7.3)	(100.0)			
悪性高血圧	1,829	163	1,992	7	619	2,618
(%)	(91.8)	(8.2)	(100.0)			
糖尿病性腎症	85,187	5,212	90,399	314	29,565	120,278
(%)	(94.2)	(5.8)	(100.0)			
SLE腎炎	1,626	86	1,712	5	510	2,227
(%)	(95.0)	(5.0)	(100.0)			
アミロイド腎	324	10	334		128	462
(%)	(97.0)	(3.0)	(100.0)			
痛風腎	783	44	827	1	247	1,075
(%)	(94.7)	(5.3)	(100.0)			
先天性代謝異常による腎不全	184	8	192		82	274
(%)	(95.8)	(4.2)	(100.0)			
腎・尿路結核	143	8	151		44	195
(%)	(94.7)	(5.3)	(100.0)			
腎・尿路結石	418	33	451		130	581
(%)	(92.7)	(7.3)	(100.0)			
腎・尿路腫瘍	657	32	689		221	910
(%)	(95.4)	(4.6)	(100.0)			
閉塞性尿路障害	538	27	565		165	730
(%)	(95.2)	(4.8)	(100.0)			
骨髄腫	202	6	208	1	82	291
(%)	(97.1)	(2.9)	(100.0)			
腎形成不全	486	22	508	2	150	660
(%)	(95.7)	(4.3)	(100.0)			
不明	20,366	1,543	21,909	86	7,902	29,897
(%)	(93.0)	(7.0)	(100.0)			
再導入	1,552	83	1,635	3	545	2,183
(%)	(94.9)	(5.1)	(100.0)			
その他	5,488	352	5,840	35	2,169	8,044
(%)	(94.0)	(6.0)	(100.0)			
合計	220,405	15,283	235,688	736	76,772	313,196
(%)	(93.5)	(6.5)	(100.0)			
記載なし	13	1	14		7	21
(%)	(92.9)	(7.1)	(100.0)			
総計	220,418	15,284	235,702	736	76,779	313,217
(%)	(93.5)	(6.5)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査