

表 861 脳出血の既往 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	47,916	3,402	51,318	143	12,013	63,474
(%)	(93.4)	(6.6)	(100.0)			
慢性腎盂腎炎	1,560	103	1,663	3	374	2,040
(%)	(93.8)	(6.2)	(100.0)			
急速進行性糸球体腎炎	1,403	58	1,461	8	302	1,771
(%)	(96.0)	(4.0)	(100.0)			
妊娠腎/妊娠中毒症	743	59	802	2	177	981
(%)	(92.6)	(7.4)	(100.0)			
その他の分類不能の腎炎	702	39	741	3	185	929
(%)	(94.7)	(5.3)	(100.0)			
多発性嚢胞腎	5,781	866	6,647	10	1,359	8,016
(%)	(87.0)	(13.0)	(100.0)			
腎硬化症	16,842	1,301	18,143	45	3,819	22,007
(%)	(92.8)	(7.2)	(100.0)			
悪性高血圧	1,411	119	1,530	6	325	1,861
(%)	(92.2)	(7.8)	(100.0)			
糖尿病性腎症	68,134	4,148	72,282	243	16,901	89,426
(%)	(94.3)	(5.7)	(100.0)			
SLE腎炎	1,178	59	1,237	4	263	1,504
(%)	(95.2)	(4.8)	(100.0)			
アミロイド腎	230	7	237		77	314
(%)	(97.0)	(3.0)	(100.0)			
痛風腎	603	38	641	1	136	778
(%)	(94.1)	(5.9)	(100.0)			
先天性代謝異常による腎不全	135	4	139		34	173
(%)	(97.1)	(2.9)	(100.0)			
腎・尿路結核	111	6	117		24	141
(%)	(94.9)	(5.1)	(100.0)			
腎・尿路結石	328	23	351		70	421
(%)	(93.4)	(6.6)	(100.0)			
腎・尿路腫瘍	532	22	554		124	678
(%)	(96.0)	(4.0)	(100.0)			
閉塞性尿路障害	392	21	413		81	494
(%)	(94.9)	(5.1)	(100.0)			
骨髄腫	142	6	148		42	190
(%)	(95.9)	(4.1)	(100.0)			
腎形成不全	316	15	331	2	67	400
(%)	(95.5)	(4.5)	(100.0)			
不明	15,590	1,215	16,805	57	4,314	21,176
(%)	(92.8)	(7.2)	(100.0)			
再導入	1,065	55	1,120	2	234	1,356
(%)	(95.1)	(4.9)	(100.0)			
その他	4,166	277	4,443	30	1,242	5,715
(%)	(93.8)	(6.2)	(100.0)			
合計	169,280	11,843	181,123	559	42,163	223,845
(%)	(93.5)	(6.5)	(100.0)			
記載なし	6	1	7			7
(%)	(85.7)	(14.3)	(100.0)			
総計	169,286	11,844	181,130	559	42,163	223,852
(%)	(93.5)	(6.5)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査