

表 962 脳梗塞の既往 原疾患別（透析患者全体）

| 原疾患           | 既往なし    | 既往あり   | 合計      | 不明  | 記載なし   | 総計      |
|---------------|---------|--------|---------|-----|--------|---------|
| 慢性糸球体腎炎       | 60,559  | 10,606 | 71,165  | 196 | 21,986 | 93,347  |
| (%)           | (85.1)  | (14.9) | (100.0) |     |        |         |
| 慢性腎盂腎炎        | 1,976   | 311    | 2,287   | 6   | 642    | 2,935   |
| (%)           | (86.4)  | (13.6) | (100.0) |     |        |         |
| 急速進行性糸球体腎炎    | 1,661   | 248    | 1,909   | 9   | 560    | 2,478   |
| (%)           | (87.0)  | (13.0) | (100.0) |     |        |         |
| 妊娠腎/妊娠中毒症     | 1,022   | 175    | 1,197   | 4   | 344    | 1,545   |
| (%)           | (85.4)  | (14.6) | (100.0) |     |        |         |
| その他の分類不能の腎炎   | 925     | 126    | 1,051   | 3   | 351    | 1,405   |
| (%)           | (88.0)  | (12.0) | (100.0) |     |        |         |
| 多発性嚢胞腎        | 7,412   | 1,348  | 8,760   | 14  | 2,482  | 11,256  |
| (%)           | (84.6)  | (15.4) | (100.0) |     |        |         |
| 腎硬化症          | 18,631  | 4,607  | 23,238  | 60  | 6,507  | 29,805  |
| (%)           | (80.2)  | (19.8) | (100.0) |     |        |         |
| 悪性高血圧         | 1,696   | 318    | 2,014   | 7   | 597    | 2,618   |
| (%)           | (84.2)  | (15.8) | (100.0) |     |        |         |
| 糖尿病性腎症        | 71,479  | 21,234 | 92,713  | 307 | 27,258 | 120,278 |
| (%)           | (77.1)  | (22.9) | (100.0) |     |        |         |
| SLE腎炎         | 1,493   | 234    | 1,727   | 5   | 495    | 2,227   |
| (%)           | (86.5)  | (13.5) | (100.0) |     |        |         |
| アミロイド腎        | 303     | 37     | 340     |     | 122    | 462     |
| (%)           | (89.1)  | (10.9) | (100.0) |     |        |         |
| 痛風腎           | 714     | 117    | 831     | 3   | 241    | 1,075   |
| (%)           | (85.9)  | (14.1) | (100.0) |     |        |         |
| 先天性代謝異常による腎不全 | 172     | 21     | 193     |     | 81     | 274     |
| (%)           | (89.1)  | (10.9) | (100.0) |     |        |         |
| 腎・尿路結核        | 131     | 22     | 153     |     | 42     | 195     |
| (%)           | (85.6)  | (14.4) | (100.0) |     |        |         |
| 腎・尿路結石        | 380     | 78     | 458     | 1   | 122    | 581     |
| (%)           | (83.0)  | (17.0) | (100.0) |     |        |         |
| 腎・尿路腫瘍        | 593     | 100    | 693     | 1   | 216    | 910     |
| (%)           | (85.6)  | (14.4) | (100.0) |     |        |         |
| 閉塞性尿路障害       | 507     | 63     | 570     |     | 160    | 730     |
| (%)           | (88.9)  | (11.1) | (100.0) |     |        |         |
| 骨髄腫           | 194     | 15     | 209     | 1   | 81     | 291     |
| (%)           | (92.8)  | (7.2)  | (100.0) |     |        |         |
| 腎形成不全         | 480     | 31     | 511     |     | 149    | 660     |
| (%)           | (93.9)  | (6.1)  | (100.0) |     |        |         |
| 不明            | 18,434  | 3,866  | 22,300  | 80  | 7,517  | 29,897  |
| (%)           | (82.7)  | (17.3) | (100.0) |     |        |         |
| 再導入           | 1,468   | 176    | 1,644   | 3   | 536    | 2,183   |
| (%)           | (89.3)  | (10.7) | (100.0) |     |        |         |
| その他           | 5,020   | 900    | 5,920   | 32  | 2,092  | 8,044   |
| (%)           | (84.8)  | (15.2) | (100.0) |     |        |         |
| 合計            | 195,250 | 44,633 | 239,883 | 732 | 72,581 | 313,196 |
| (%)           | (81.4)  | (18.6) | (100.0) |     |        |         |
| 記載なし          | 12      | 2      | 14      |     | 7      | 21      |
| (%)           | (85.7)  | (14.3) | (100.0) |     |        |         |
| 総計            | 195,262 | 44,635 | 239,897 | 732 | 72,588 | 313,217 |
| (%)           | (81.4)  | (18.6) | (100.0) |     |        |         |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査