

表 990 脳梗塞の既往 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	43,981	7,872	51,853	142	11,479	63,474
(%)	(84.8)	(15.2)	(100.0)			
慢性腎盂腎炎	1,454	233	1,687	5	348	2,040
(%)	(86.2)	(13.8)	(100.0)			
急速進行性糸球体腎炎	1,287	189	1,476	8	287	1,771
(%)	(87.2)	(12.8)	(100.0)			
妊娠腎/妊娠中毒症	692	114	806	3	172	981
(%)	(85.9)	(14.1)	(100.0)			
その他の分類不能の腎炎	645	100	745	3	181	929
(%)	(86.6)	(13.4)	(100.0)			
多発性嚢胞腎	5,666	992	6,658	11	1,347	8,016
(%)	(85.1)	(14.9)	(100.0)			
腎硬化症	14,769	3,631	18,400	42	3,565	22,007
(%)	(80.3)	(19.7)	(100.0)			
悪性高血圧	1,287	257	1,544	7	310	1,861
(%)	(83.4)	(16.6)	(100.0)			
糖尿病性腎症	57,045	16,819	73,864	239	15,323	89,426
(%)	(77.2)	(22.8)	(100.0)			
SLE腎炎	1,073	175	1,248	4	252	1,504
(%)	(86.0)	(14.0)	(100.0)			
アミロイド腎	217	24	241		73	314
(%)	(90.0)	(10.0)	(100.0)			
痛風腎	550	95	645	3	130	778
(%)	(85.3)	(14.7)	(100.0)			
先天性代謝異常による腎不全	124	16	140		33	173
(%)	(88.6)	(11.4)	(100.0)			
腎・尿路結核	102	16	118		23	141
(%)	(86.4)	(13.6)	(100.0)			
腎・尿路結石	299	56	355		66	421
(%)	(84.2)	(15.8)	(100.0)			
腎・尿路腫瘍	480	75	555	1	122	678
(%)	(86.5)	(13.5)	(100.0)			
閉塞性尿路障害	372	42	414		80	494
(%)	(89.9)	(10.1)	(100.0)			
骨髄腫	137	12	149		41	190
(%)	(91.9)	(8.1)	(100.0)			
腎形成不全	312	21	333		67	400
(%)	(93.7)	(6.3)	(100.0)			
不明	14,101	2,965	17,066	49	4,061	21,176
(%)	(82.6)	(17.4)	(100.0)			
再導入	1,004	120	1,124	2	230	1,356
(%)	(89.3)	(10.7)	(100.0)			
その他	3,807	695	4,502	26	1,187	5,715
(%)	(84.6)	(15.4)	(100.0)			
合計	149,404	34,519	183,923	545	39,377	223,845
(%)	(81.2)	(18.8)	(100.0)			
記載なし	6	1	7			7
(%)	(85.7)	(14.3)	(100.0)			
総計	149,410	34,520	183,930	545	39,377	223,852
(%)	(81.2)	(18.8)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査