

表 1058 脳梗塞の既往 原疾患別（腹膜透析患者）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	1,743	154	1,897	5	928	2,830
(%)	(91.9)	(8.1)	(100.0)			
慢性腎盂腎炎	69	8	77		20	97
(%)	(89.6)	(10.4)	(100.0)			
急速進行性糸球体腎炎	34	6	40		26	66
(%)	(85.0)	(15.0)	(100.0)			
妊娠腎/妊娠中毒症	22	2	24		3	27
(%)	(91.7)	(8.3)	(100.0)			
その他の分類不能の腎炎	53	1	54		24	78
(%)	(98.1)	(1.9)	(100.0)			
多発性嚢胞腎	153	17	170		93	263
(%)	(90.0)	(10.0)	(100.0)			
腎硬化症	627	109	736	1	358	1,095
(%)	(85.2)	(14.8)	(100.0)			
悪性高血圧	62	5	67		40	107
(%)	(92.5)	(7.5)	(100.0)			
糖尿病性腎症	1,538	330	1,868	8	923	2,799
(%)	(82.3)	(17.7)	(100.0)			
SLE腎炎	28	6	34		12	46
(%)	(82.4)	(17.6)	(100.0)			
アミロイド腎	9	1	10		5	15
(%)	(90.0)	(10.0)	(100.0)			
痛風腎	26		26		6	32
(%)	(100.0)		(100.0)			
先天性代謝異常による腎不全	8	3	11		6	17
(%)	(72.7)	(27.3)	(100.0)			
腎・尿路結核					2	2
(%)						
腎・尿路結石	11	5	16		5	21
(%)	(68.8)	(31.3)	(100.0)			
腎・尿路腫瘍	10	1	11		3	14
(%)	(90.9)	(9.1)	(100.0)			
閉塞性尿路障害	14	1	15		6	21
(%)	(93.3)	(6.7)	(100.0)			
骨髄腫	1	2	3		4	7
(%)	(33.3)	(66.7)	(100.0)			
腎形成不全	47	2	49		23	72
(%)	(95.9)	(4.1)	(100.0)			
不明	488	63	551	7	284	842
(%)	(88.6)	(11.4)	(100.0)			
再導入	27	2	29		26	55
(%)	(93.1)	(6.9)	(100.0)			
その他	152	13	165	1	97	263
(%)	(92.1)	(7.9)	(100.0)			
合計	5,122	731	5,853	22	2,894	8,769
(%)	(87.5)	(12.5)	(100.0)			
記載なし						
(%)						
総計	5,122	731	5,853	22	2,894	8,769
(%)	(87.5)	(12.5)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査