

表 1088 四肢切断の有無 原疾患別（透析患者全体）

| 原疾患           | 切断なし    | 切断あり  | 合計      | 不明  | 記載なし   | 総計      |
|---------------|---------|-------|---------|-----|--------|---------|
| 慢性糸球体腎炎       | 69,521  | 1,038 | 70,559  | 73  | 22,715 | 93,347  |
| (%)           | (98.5)  | (1.5) | (100.0) |     |        |         |
| 慢性腎盂腎炎        | 2,234   | 46    | 2,280   | 1   | 654    | 2,935   |
| (%)           | (98.0)  | (2.0) | (100.0) |     |        |         |
| 急速進行性糸球体腎炎    | 1,874   | 25    | 1,899   | 3   | 576    | 2,478   |
| (%)           | (98.7)  | (1.3) | (100.0) |     |        |         |
| 妊娠腎/妊娠中毒症     | 1,174   | 16    | 1,190   | 3   | 352    | 1,545   |
| (%)           | (98.7)  | (1.3) | (100.0) |     |        |         |
| その他の分類不能の腎炎   | 1,030   | 9     | 1,039   | 3   | 363    | 1,405   |
| (%)           | (99.1)  | (0.9) | (100.0) |     |        |         |
| 多発性嚢胞腎        | 8,614   | 68    | 8,682   | 5   | 2,569  | 11,256  |
| (%)           | (99.2)  | (0.8) | (100.0) |     |        |         |
| 腎硬化症          | 22,550  | 334   | 22,884  | 20  | 6,901  | 29,805  |
| (%)           | (98.5)  | (1.5) | (100.0) |     |        |         |
| 悪性高血圧         | 1,961   | 30    | 1,991   | 2   | 625    | 2,618   |
| (%)           | (98.5)  | (1.5) | (100.0) |     |        |         |
| 糖尿病性腎症        | 84,777  | 6,894 | 91,671  | 103 | 28,504 | 120,278 |
| (%)           | (92.5)  | (7.5) | (100.0) |     |        |         |
| SLE腎炎         | 1,674   | 47    | 1,721   | 2   | 504    | 2,227   |
| (%)           | (97.3)  | (2.7) | (100.0) |     |        |         |
| アミロイド腎        | 330     | 2     | 332     |     | 130    | 462     |
| (%)           | (99.4)  | (0.6) | (100.0) |     |        |         |
| 痛風腎           | 824     | 11    | 835     | 1   | 239    | 1,075   |
| (%)           | (98.7)  | (1.3) | (100.0) |     |        |         |
| 先天性代謝異常による腎不全 | 185     | 5     | 190     |     | 84     | 274     |
| (%)           | (97.4)  | (2.6) | (100.0) |     |        |         |
| 腎・尿路結核        | 144     | 3     | 147     |     | 48     | 195     |
| (%)           | (98.0)  | (2.0) | (100.0) |     |        |         |
| 腎・尿路結石        | 446     | 10    | 456     |     | 125    | 581     |
| (%)           | (97.8)  | (2.2) | (100.0) |     |        |         |
| 腎・尿路腫瘍        | 688     | 9     | 697     |     | 213    | 910     |
| (%)           | (98.7)  | (1.3) | (100.0) |     |        |         |
| 閉塞性尿路障害       | 558     | 16    | 574     |     | 156    | 730     |
| (%)           | (97.2)  | (2.8) | (100.0) |     |        |         |
| 骨髄腫           | 209     | 1     | 210     |     | 81     | 291     |
| (%)           | (99.5)  | (0.5) | (100.0) |     |        |         |
| 腎形成不全         | 501     | 9     | 510     |     | 150    | 660     |
| (%)           | (98.2)  | (1.8) | (100.0) |     |        |         |
| 不明            | 21,525  | 457   | 21,982  | 17  | 7,898  | 29,897  |
| (%)           | (97.9)  | (2.1) | (100.0) |     |        |         |
| 再導入           | 1,600   | 44    | 1,644   |     | 539    | 2,183   |
| (%)           | (97.3)  | (2.7) | (100.0) |     |        |         |
| その他           | 5,781   | 133   | 5,914   | 10  | 2,120  | 8,044   |
| (%)           | (97.8)  | (2.2) | (100.0) |     |        |         |
| 合計            | 228,200 | 9,207 | 237,407 | 243 | 75,546 | 313,196 |
| (%)           | (96.1)  | (3.9) | (100.0) |     |        |         |
| 記載なし          | 13      | 1     | 14      |     | 7      | 21      |
| (%)           | (92.9)  | (7.1) | (100.0) |     |        |         |
| 総計            | 228,213 | 9,208 | 237,421 | 243 | 75,553 | 313,217 |
| (%)           | (96.1)  | (3.9) | (100.0) |     |        |         |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査