

表 1115 四肢切断の有無 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	切断なし	切断あり	合計	不明	記載なし	総計
慢性糸球体腎炎	50,691	797	51,488	63	11,923	63,474
(%)	(98.5)	(1.5)	(100.0)			
慢性腎盂腎炎	1,653	29	1,682	1	357	2,040
(%)	(98.3)	(1.7)	(100.0)			
急速進行性糸球体腎炎	1,451	22	1,473	3	295	1,771
(%)	(98.5)	(1.5)	(100.0)			
妊娠腎/妊娠中毒症	796	11	807	3	171	981
(%)	(98.6)	(1.4)	(100.0)			
その他の分類不能の腎炎	729	8	737	3	189	929
(%)	(98.9)	(1.1)	(100.0)			
多発性嚢胞腎	6,569	50	6,619	4	1,393	8,016
(%)	(99.2)	(0.8)	(100.0)			
腎硬化症	17,925	263	18,188	18	3,801	22,007
(%)	(98.6)	(1.4)	(100.0)			
悪性高血圧	1,508	24	1,532	2	327	1,861
(%)	(98.4)	(1.6)	(100.0)			
糖尿病性腎症	67,740	5,479	73,219	94	16,113	89,426
(%)	(92.5)	(7.5)	(100.0)			
SLE腎炎	1,207	42	1,249	1	254	1,504
(%)	(96.6)	(3.4)	(100.0)			
アミロイド腎	233	2	235		79	314
(%)	(99.1)	(0.9)	(100.0)			
痛風腎	639	8	647	1	130	778
(%)	(98.8)	(1.2)	(100.0)			
先天性代謝異常による腎不全	135	3	138		35	173
(%)	(97.8)	(2.2)	(100.0)			
腎・尿路結核	112	2	114		27	141
(%)	(98.2)	(1.8)	(100.0)			
腎・尿路結石	351	5	356		65	421
(%)	(98.6)	(1.4)	(100.0)			
腎・尿路腫瘍	555	7	562		116	678
(%)	(98.8)	(1.2)	(100.0)			
閉塞性尿路障害	403	16	419		75	494
(%)	(96.2)	(3.8)	(100.0)			
骨髄腫	148	1	149		41	190
(%)	(99.3)	(0.7)	(100.0)			
腎形成不全	327	6	333		67	400
(%)	(98.2)	(1.8)	(100.0)			
不明	16,484	362	16,846	11	4,319	21,176
(%)	(97.9)	(2.1)	(100.0)			
再導入	1,096	27	1,123		233	1,356
(%)	(97.6)	(2.4)	(100.0)			
その他	4,399	103	4,502	10	1,203	5,715
(%)	(97.7)	(2.3)	(100.0)			
合計	175,151	7,267	182,418	214	41,213	223,845
(%)	(96.0)	(4.0)	(100.0)			
記載なし	6	1	7			7
(%)	(85.7)	(14.3)	(100.0)			
総計	175,157	7,268	182,425	214	41,213	223,852
(%)	(96.0)	(4.0)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査