

表 1210 大腿骨近位部骨折の既往 原疾患別（透析患者全体）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	67,218	2,957	70,175	112	23,060	93,347
(%)	(95.8)	(4.2)	(100.0)			
慢性腎盂腎炎	2,140	124	2,264	3	668	2,935
(%)	(94.5)	(5.5)	(100.0)			
急速進行性糸球体腎炎	1,808	83	1,891	1	586	2,478
(%)	(95.6)	(4.4)	(100.0)			
妊娠腎/妊娠中毒症	1,103	75	1,178	5	362	1,545
(%)	(93.6)	(6.4)	(100.0)			
その他の分類不能の腎炎	981	52	1,033	3	369	1,405
(%)	(95.0)	(5.0)	(100.0)			
多発性嚢胞腎	8,347	272	8,619	8	2,629	11,256
(%)	(96.8)	(3.2)	(100.0)			
腎硬化症	21,619	1,142	22,761	29	7,015	29,805
(%)	(95.0)	(5.0)	(100.0)			
悪性高血圧	1,921	65	1,986	3	629	2,618
(%)	(96.7)	(3.3)	(100.0)			
糖尿病性腎症	85,773	4,548	90,321	171	29,786	120,278
(%)	(95.0)	(5.0)	(100.0)			
SLE腎炎	1,603	115	1,718	5	504	2,227
(%)	(93.3)	(6.7)	(100.0)			
アミロイド腎	310	25	335		127	462
(%)	(92.5)	(7.5)	(100.0)			
痛風腎	812	21	833	2	240	1,075
(%)	(97.5)	(2.5)	(100.0)			
先天性代謝異常による腎不全	181	9	190		84	274
(%)	(95.3)	(4.7)	(100.0)			
腎・尿路結核	138	11	149		46	195
(%)	(92.6)	(7.4)	(100.0)			
腎・尿路結石	437	17	454		127	581
(%)	(96.3)	(3.7)	(100.0)			
腎・尿路腫瘍	680	15	695		215	910
(%)	(97.8)	(2.2)	(100.0)			
閉塞性尿路障害	549	18	567		163	730
(%)	(96.8)	(3.2)	(100.0)			
骨髄腫	202	8	210	1	80	291
(%)	(96.2)	(3.8)	(100.0)			
腎形成不全	495	9	504		156	660
(%)	(98.2)	(1.8)	(100.0)			
不明	20,730	1,095	21,825	66	8,006	29,897
(%)	(95.0)	(5.0)	(100.0)			
再導入	1,578	63	1,641	2	540	2,183
(%)	(96.2)	(3.8)	(100.0)			
その他	5,533	287	5,820	31	2,193	8,044
(%)	(95.1)	(4.9)	(100.0)			
合計	224,158	11,011	235,169	442	77,585	313,196
(%)	(95.3)	(4.7)	(100.0)			
記載なし	14		14		7	21
(%)	(100.0)		(100.0)			
総計	224,172	11,011	235,183	442	77,592	313,217
(%)	(95.3)	(4.7)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査