

表 1240 大腿骨近位部骨折の既往 原疾患別（施設血液透析患者 透析回数週3回）

| 原疾患           | 既往なし    | 既往あり  | 合計      | 不明  | 記載なし   | 総計      |
|---------------|---------|-------|---------|-----|--------|---------|
| 慢性糸球体腎炎       | 49,034  | 2,240 | 51,274  | 93  | 12,107 | 63,474  |
| (%)           | (95.6)  | (4.4) | (100.0) |     |        |         |
| 慢性腎盂腎炎        | 1,578   | 96    | 1,674   | 3   | 363    | 2,040   |
| (%)           | (94.3)  | (5.7) | (100.0) |     |        |         |
| 急速進行性糸球体腎炎    | 1,399   | 67    | 1,466   | 1   | 304    | 1,771   |
| (%)           | (95.4)  | (4.6) | (100.0) |     |        |         |
| 妊娠腎/妊娠中毒症     | 747     | 54    | 801     | 3   | 177    | 981     |
| (%)           | (93.3)  | (6.7) | (100.0) |     |        |         |
| その他の分類不能の腎炎   | 694     | 41    | 735     | 3   | 191    | 929     |
| (%)           | (94.4)  | (5.6) | (100.0) |     |        |         |
| 多発性嚢胞腎        | 6,372   | 203   | 6,575   | 6   | 1,435  | 8,016   |
| (%)           | (96.9)  | (3.1) | (100.0) |     |        |         |
| 腎硬化症          | 17,193  | 910   | 18,103  | 19  | 3,885  | 22,007  |
| (%)           | (95.0)  | (5.0) | (100.0) |     |        |         |
| 悪性高血圧         | 1,480   | 50    | 1,530   | 3   | 328    | 1,861   |
| (%)           | (96.7)  | (3.3) | (100.0) |     |        |         |
| 糖尿病性腎症        | 68,597  | 3,728 | 72,325  | 149 | 16,952 | 89,426  |
| (%)           | (94.8)  | (5.2) | (100.0) |     |        |         |
| SLE腎炎         | 1,163   | 83    | 1,246   | 4   | 254    | 1,504   |
| (%)           | (93.3)  | (6.7) | (100.0) |     |        |         |
| アミロイド腎        | 218     | 20    | 238     |     | 76     | 314     |
| (%)           | (91.6)  | (8.4) | (100.0) |     |        |         |
| 痛風腎           | 632     | 14    | 646     | 2   | 130    | 778     |
| (%)           | (97.8)  | (2.2) | (100.0) |     |        |         |
| 先天性代謝異常による腎不全 | 131     | 8     | 139     |     | 34     | 173     |
| (%)           | (94.2)  | (5.8) | (100.0) |     |        |         |
| 腎・尿路結核        | 104     | 11    | 115     |     | 26     | 141     |
| (%)           | (90.4)  | (9.6) | (100.0) |     |        |         |
| 腎・尿路結石        | 344     | 9     | 353     |     | 68     | 421     |
| (%)           | (97.5)  | (2.5) | (100.0) |     |        |         |
| 腎・尿路腫瘍        | 546     | 15    | 561     |     | 117    | 678     |
| (%)           | (97.3)  | (2.7) | (100.0) |     |        |         |
| 閉塞性尿路障害       | 401     | 15    | 416     |     | 78     | 494     |
| (%)           | (96.4)  | (3.6) | (100.0) |     |        |         |
| 骨髄腫           | 144     | 5     | 149     |     | 41     | 190     |
| (%)           | (96.6)  | (3.4) | (100.0) |     |        |         |
| 腎形成不全         | 322     | 7     | 329     |     | 71     | 400     |
| (%)           | (97.9)  | (2.1) | (100.0) |     |        |         |
| 不明            | 15,871  | 875   | 16,746  | 52  | 4,378  | 21,176  |
| (%)           | (94.8)  | (5.2) | (100.0) |     |        |         |
| 再導入           | 1,083   | 37    | 1,120   | 1   | 235    | 1,356   |
| (%)           | (96.7)  | (3.3) | (100.0) |     |        |         |
| その他           | 4,218   | 219   | 4,437   | 25  | 1,253  | 5,715   |
| (%)           | (95.1)  | (4.9) | (100.0) |     |        |         |
| 合計            | 172,271 | 8,707 | 180,978 | 364 | 42,503 | 223,845 |
| (%)           | (95.2)  | (4.8) | (100.0) |     |        |         |
| 記載なし          | 7       |       | 7       |     |        | 7       |
| (%)           | (100.0) |       | (100.0) |     |        |         |
| 総計            | 172,278 | 8,707 | 180,985 | 364 | 42,503 | 223,852 |
| (%)           | (95.2)  | (4.8) | (100.0) |     |        |         |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査