

表 1403 降圧薬使用の有無 原疾患別（透析患者全体）

原疾患	なし	あり	合計	不明	記載なし	総計
慢性糸球体腎炎	27,798	46,193	73,991	229	19,127	93,347
(%)	(37.6)	(62.4)	(100.0)			
慢性腎盂腎炎	978	1,398	2,376	9	550	2,935
(%)	(41.2)	(58.8)	(100.0)			
急速進行性糸球体腎炎	760	1,231	1,991	12	475	2,478
(%)	(38.2)	(61.8)	(100.0)			
妊娠腎/妊娠中毒症	491	755	1,246	8	291	1,545
(%)	(39.4)	(60.6)	(100.0)			
その他の分類不能の腎炎	424	669	1,093	6	306	1,405
(%)	(38.8)	(61.2)	(100.0)			
多発性嚢胞腎	3,346	5,711	9,057	33	2,166	11,256
(%)	(36.9)	(63.1)	(100.0)			
腎硬化症	7,798	16,125	23,923	70	5,812	29,805
(%)	(32.6)	(67.4)	(100.0)			
悪性高血圧	556	1,567	2,123	11	484	2,618
(%)	(26.2)	(73.8)	(100.0)			
糖尿病性腎症	25,539	69,896	95,435	418	24,425	120,278
(%)	(26.8)	(73.2)	(100.0)			
SLE腎炎	702	1,095	1,797	5	425	2,227
(%)	(39.1)	(60.9)	(100.0)			
アミロイド腎	147	209	356	1	105	462
(%)	(41.3)	(58.7)	(100.0)			
痛風腎	344	538	882	3	190	1,075
(%)	(39.0)	(61.0)	(100.0)			
先天性代謝異常による腎不全	95	111	206	1	67	274
(%)	(46.1)	(53.9)	(100.0)			
腎・尿路結核	82	73	155	1	39	195
(%)	(52.9)	(47.1)	(100.0)			
腎・尿路結石	198	268	466	1	114	581
(%)	(42.5)	(57.5)	(100.0)			
腎・尿路腫瘍	271	454	725	2	183	910
(%)	(37.4)	(62.6)	(100.0)			
閉塞性尿路障害	265	322	587		143	730
(%)	(45.1)	(54.9)	(100.0)			
骨髄腫	132	91	223	1	67	291
(%)	(59.2)	(40.8)	(100.0)			
腎形成不全	230	299	529	3	128	660
(%)	(43.5)	(56.5)	(100.0)			
不明	8,633	14,375	23,008	163	6,726	29,897
(%)	(37.5)	(62.5)	(100.0)			
再導入	676	1,054	1,730	1	452	2,183
(%)	(39.1)	(60.9)	(100.0)			
その他	2,420	3,759	6,179	61	1,804	8,044
(%)	(39.2)	(60.8)	(100.0)			
合計	81,885	166,193	248,078	1,039	64,079	313,196
(%)	(33.0)	(67.0)	(100.0)			
記載なし	4	4	8		13	21
(%)	(50.0)	(50.0)	(100.0)			
総計	81,889	166,197	248,086	1,039	64,092	313,217
(%)	(33.0)	(67.0)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査