

表 1427 降圧薬使用の有無 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	なし	あり	合計	不明	記載なし	総計
慢性糸球体腎炎	20,168	33,896	54,064	178	9,232	63,474
(%)	(37.3)	(62.7)	(100.0)			
慢性腎盂腎炎	718	1,040	1,758	8	274	2,040
(%)	(40.8)	(59.2)	(100.0)			
急速進行性糸球体腎炎	569	965	1,534	11	226	1,771
(%)	(37.1)	(62.9)	(100.0)			
妊娠腎/妊娠中毒症	324	514	838	7	136	981
(%)	(38.7)	(61.3)	(100.0)			
その他の分類不能の腎炎	304	477	781	5	143	929
(%)	(38.9)	(61.1)	(100.0)			
多発性嚢胞腎	2,528	4,380	6,908	27	1,081	8,016
(%)	(36.6)	(63.4)	(100.0)			
腎硬化症	6,253	12,759	19,012	55	2,940	22,007
(%)	(32.9)	(67.1)	(100.0)			
悪性高血圧	436	1,189	1,625	10	226	1,861
(%)	(26.8)	(73.2)	(100.0)			
糖尿病性腎症	20,402	55,949	76,351	357	12,718	89,426
(%)	(26.7)	(73.3)	(100.0)			
SLE腎炎	504	808	1,312	3	189	1,504
(%)	(38.4)	(61.6)	(100.0)			
アミロイド腎	104	154	258	1	55	314
(%)	(40.3)	(59.7)	(100.0)			
痛風腎	266	416	682	2	94	778
(%)	(39.0)	(61.0)	(100.0)			
先天性代謝異常による腎不全	71	74	145	1	27	173
(%)	(49.0)	(51.0)	(100.0)			
腎・尿路結核	60	58	118	1	22	141
(%)	(50.8)	(49.2)	(100.0)			
腎・尿路結石	155	210	365	1	55	421
(%)	(42.5)	(57.5)	(100.0)			
腎・尿路腫瘍	228	363	591	1	86	678
(%)	(38.6)	(61.4)	(100.0)			
閉塞性尿路障害	196	238	434		60	494
(%)	(45.2)	(54.8)	(100.0)			
骨髄腫	90	66	156	1	33	190
(%)	(57.7)	(42.3)	(100.0)			
腎形成不全	151	202	353		47	400
(%)	(42.8)	(57.2)	(100.0)			
不明	6,601	11,055	17,656	126	3,394	21,176
(%)	(37.4)	(62.6)	(100.0)			
再導入	456	713	1,169		187	1,356
(%)	(39.0)	(61.0)	(100.0)			
その他	1,840	2,873	4,713	53	949	5,715
(%)	(39.0)	(61.0)	(100.0)			
合計	62,424	128,399	190,823	848	32,174	223,845
(%)	(32.7)	(67.3)	(100.0)			
記載なし	3		3		4	7
(%)	(100.0)		(100.0)			
総計	62,427	128,399	190,826	848	32,178	223,852
(%)	(32.7)	(67.3)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査