

表 1458 降圧薬使用の有無 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	なし	あり	合計	不明	記載なし	総計
慢性糸球体腎炎	5,869	9,019	14,888	33	2,541	17,462
(%)	(39.4)	(60.6)	(100.0)			
慢性腎盂腎炎	182	235	417	1	66	484
(%)	(43.6)	(56.4)	(100.0)			
急速進行性糸球体腎炎	130	178	308	1	44	353
(%)	(42.2)	(57.8)	(100.0)			
妊娠腎/妊娠中毒症	139	189	328	1	72	401
(%)	(42.4)	(57.6)	(100.0)			
その他の分類不能の腎炎	89	121	210	1	50	261
(%)	(42.4)	(57.6)	(100.0)			
多発性嚢胞腎	636	964	1,600	3	277	1,880
(%)	(39.8)	(60.3)	(100.0)			
腎硬化症	1,085	2,144	3,229	5	586	3,820
(%)	(33.6)	(66.4)	(100.0)			
悪性高血圧	93	253	346		45	391
(%)	(26.9)	(73.1)	(100.0)			
糖尿病性腎症	3,861	10,432	14,293	29	2,638	16,960
(%)	(27.0)	(73.0)	(100.0)			
SLE腎炎	154	219	373	2	51	426
(%)	(41.3)	(58.7)	(100.0)			
アミロイド腎	28	37	65		9	74
(%)	(43.1)	(56.9)	(100.0)			
痛風腎	65	83	148		18	166
(%)	(43.9)	(56.1)	(100.0)			
先天性代謝異常による腎不全	13	28	41		8	49
(%)	(31.7)	(68.3)	(100.0)			
腎・尿路結核	14	13	27		4	31
(%)	(51.9)	(48.1)	(100.0)			
腎・尿路結石	25	40	65		14	79
(%)	(38.5)	(61.5)	(100.0)			
腎・尿路腫瘍	31	71	102		17	119
(%)	(30.4)	(69.6)	(100.0)			
閉塞性尿路障害	36	49	85		19	104
(%)	(42.4)	(57.6)	(100.0)			
骨髄腫	15	14	29		4	33
(%)	(51.7)	(48.3)	(100.0)			
腎形成不全	54	57	111		25	136
(%)	(48.6)	(51.4)	(100.0)			
不明	1,431	2,338	3,769	15	919	4,703
(%)	(38.0)	(62.0)	(100.0)			
再導入	163	261	424		59	483
(%)	(38.4)	(61.6)	(100.0)			
その他	372	586	958	6	196	1,160
(%)	(38.8)	(61.2)	(100.0)			
合計	14,485	27,331	41,816	97	7,662	49,575
(%)	(34.6)	(65.4)	(100.0)			
記載なし	1	3	4		2	6
(%)	(25.0)	(75.0)	(100.0)			
総計	14,486	27,334	41,820	97	7,664	49,581
(%)	(34.6)	(65.4)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査