

表 1489 降圧薬使用の有無 原疾患別（腹膜透析患者）

原疾患	なし	あり	合計	不明	記載なし	総計
慢性糸球体腎炎	393	1,577	1,970	10	850	2,830
(%)	(19.9)	(80.1)	(100.0)			
慢性腎盂腎炎	22	58	80		17	97
(%)	(27.5)	(72.5)	(100.0)			
急速進行性糸球体腎炎	10	34	44		22	66
(%)	(22.7)	(77.3)	(100.0)			
妊娠腎/妊娠中毒症	4	20	24		3	27
(%)	(16.7)	(83.3)	(100.0)			
その他の分類不能の腎炎	10	45	55		23	78
(%)	(18.2)	(81.8)	(100.0)			
多発性嚢胞腎	48	128	176		87	263
(%)	(27.3)	(72.7)	(100.0)			
腎硬化症	143	602	745	2	348	1,095
(%)	(19.2)	(80.8)	(100.0)			
悪性高血圧	12	58	70		37	107
(%)	(17.1)	(82.9)	(100.0)			
糖尿病性腎症	375	1,493	1,868	13	918	2,799
(%)	(20.1)	(79.9)	(100.0)			
SLE腎炎	9	22	31		15	46
(%)	(29.0)	(71.0)	(100.0)			
アミロイド腎	3	7	10		5	15
(%)	(30.0)	(70.0)	(100.0)			
痛風腎	7	17	24		8	32
(%)	(29.2)	(70.8)	(100.0)			
先天性代謝異常による腎不全	6	5	11		6	17
(%)	(54.5)	(45.5)	(100.0)			
腎・尿路結核					2	2
(%)						
腎・尿路結石	7	8	15		6	21
(%)	(46.7)	(53.3)	(100.0)			
腎・尿路腫瘍	4	6	10	1	3	14
(%)	(40.0)	(60.0)	(100.0)			
閉塞性尿路障害	4	10	14		7	21
(%)	(28.6)	(71.4)	(100.0)			
骨髄腫	3		3		4	7
(%)	(100.0)		(100.0)			
腎形成不全	17	31	48	3	21	72
(%)	(35.4)	(64.6)	(100.0)			
不明	164	403	567	6	269	842
(%)	(28.9)	(71.1)	(100.0)			
再導入	4	27	31		24	55
(%)	(12.9)	(87.1)	(100.0)			
その他	50	127	177	1	85	263
(%)	(28.2)	(71.8)	(100.0)			
合計	1,295	4,678	5,973	36	2,760	8,769
(%)	(21.7)	(78.3)	(100.0)			
記載なし						
(%)						
総計	1,295	4,678	5,973	36	2,760	8,769
(%)	(21.7)	(78.3)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査