

表 1703 腎移植の回数 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
施設血液透析	180,595	1,847	92	3	3	182,540	508	65,677	248,725
(%)	(98.9)	(1.0)	(0.1)	(0.0)	(0.0)	(100.0)			
Off-line HDF	3,795	103	3			3,901	9	1,422	5,332
(%)	(97.3)	(2.6)	(0.1)			(100.0)			
On-line HDF	30,868	599	35	3	1	31,506	126	12,895	44,527
(%)	(98.0)	(1.9)	(0.1)	(0.0)	(0.0)	(100.0)			
Push/Pull HDF	67					67		43	110
(%)	(100.0)					(100.0)			
ハイオフィльтраーション	199	5				204	1	62	267
(%)	(97.5)	(2.5)				(100.0)			
IHDF	2,636	24	3			2,663		877	3,540
(%)	(99.0)	(0.9)	(0.1)			(100.0)			
血液濾過	11					11		5	16
(%)	(100.0)					(100.0)			
血液吸着透析 (リクセル等使用)	1,050	66	5	1		1,122		254	1,376
(%)	(93.6)	(5.9)	(0.4)	(0.1)		(100.0)			
在宅血液透析	372	41	4			417	5	133	555
(%)	(89.2)	(9.8)	(1.0)			(100.0)			
腹膜透析 (CAPD)	3,208	26		1		3,235	5	1,662	4,902
(%)	(99.2)	(0.8)		(0.0)		(100.0)			
腹膜透析 (APD)	1,835	13				1,848	6	921	2,775
(%)	(99.3)	(0.7)				(100.0)			
腹膜透析 (CCPD)	772	10				782	3	307	1,092
(%)	(98.7)	(1.3)				(100.0)			
合計	225,408	2,734	142	8	4	228,296	663	84,258	313,217
(%)	(98.7)	(1.2)	(0.1)	(0.0)	(0.0)	(100.0)			
記載なし (%)									
総計	225,408	2,734	142	8	4	228,296	663	84,258	313,217
(%)	(98.7)	(1.2)	(0.1)	(0.0)	(0.0)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査