

表 1708 腎移植の回数 原疾患別（透析患者全体）

原疾患	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
慢性糸球体腎炎	67,188	947	56	3		68,194	175	24,978	93,347
(%)	(98.5)	(1.4)	(0.1)	(0.0)		(100.0)			
慢性腎盂腎炎	2,161	25	2			2,188	8	739	2,935
(%)	(98.8)	(1.1)	(0.1)			(100.0)			
急速進行性糸球体腎炎	1,800	15				1,815	1	662	2,478
(%)	(99.2)	(0.8)				(100.0)			
妊娠腎/妊娠中毒症	1,122	16		1		1,139	5	401	1,545
(%)	(98.5)	(1.4)		(0.1)		(100.0)			
その他の分類不能の腎炎	1,001	20	1			1,022	5	378	1,405
(%)	(97.9)	(2.0)	(0.1)			(100.0)			
多発性嚢胞腎	8,254	31	2			8,287	12	2,957	11,256
(%)	(99.6)	(0.4)	(0.0)			(100.0)			
腎硬化症	22,037	44	4			22,085	43	7,677	29,805
(%)	(99.8)	(0.2)	(0.0)			(100.0)			
悪性高血圧	1,905	6				1,911	8	699	2,618
(%)	(99.7)	(0.3)				(100.0)			
糖尿病性腎症	87,359	157	10		4	87,530	237	32,511	120,278
(%)	(99.8)	(0.2)	(0.0)		(0.0)	(100.0)			
SLE腎炎	1,619	20				1,639	7	581	2,227
(%)	(98.8)	(1.2)				(100.0)			
アミロイド腎	331					331		131	462
(%)	(100.0)					(100.0)			
痛風腎	800	3				803	2	270	1,075
(%)	(99.6)	(0.4)				(100.0)			
先天性代謝異常による腎不全	180	17				197		77	274
(%)	(91.4)	(8.6)				(100.0)			
腎・尿路結核	145	1				146		49	195
(%)	(99.3)	(0.7)				(100.0)			
腎・尿路結石	439	1				440		141	581
(%)	(99.8)	(0.2)				(100.0)			
腎・尿路腫瘍	669	4				673		237	910
(%)	(99.4)	(0.6)				(100.0)			
閉塞性尿路障害	540	8				548	1	181	730
(%)	(98.5)	(1.5)				(100.0)			
骨髄腫	212					212	1	78	291
(%)	(100.0)					(100.0)			
腎形成不全	459	37	1			497		163	660
(%)	(92.4)	(7.4)	(0.2)			(100.0)			
不明	21,086	170	10			21,266	136	8,495	29,897
(%)	(99.2)	(0.8)	(0.0)			(100.0)			
再導入	587	1,118	51	4		1,760	2	421	2,183
(%)	(33.4)	(63.5)	(2.9)	(0.2)		(100.0)			
その他	5,500	94	5			5,599	20	2,425	8,044
(%)	(98.2)	(1.7)	(0.1)			(100.0)			
合計	225,394	2,734	142	8	4	228,282	663	84,251	313,196
(%)	(98.7)	(1.2)	(0.1)	(0.0)	(0.0)	(100.0)			
記載なし	14					14		7	21
(%)	(100.0)					(100.0)			
総計	225,408	2,734	142	8	4	228,296	663	84,258	313,217
(%)	(98.7)	(1.2)	(0.1)	(0.0)	(0.0)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査