

表 1729 腎移植の回数 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
慢性糸球体腎炎	49,522	569	36	1		50,128	131	13,215	63,474
(%)	(98.8)	(1.1)	(0.1)	(0.0)		(100.0)			
慢性腎盂腎炎	1,612	19				1,631	6	403	2,040
(%)	(98.8)	(1.2)				(100.0)			
急速進行性糸球体腎炎	1,409	8				1,417	1	353	1,771
(%)	(99.4)	(0.6)				(100.0)			
妊娠腎/妊娠中毒症	754	8				762	4	215	981
(%)	(99.0)	(1.0)				(100.0)			
その他の分類不能の腎炎	707	13				720	2	207	929
(%)	(98.2)	(1.8)				(100.0)			
多発性嚢胞腎	6,376	21	2			6,399	8	1,609	8,016
(%)	(99.6)	(0.3)	(0.0)			(100.0)			
腎硬化症	17,573	27	2			17,602	34	4,371	22,007
(%)	(99.8)	(0.2)	(0.0)			(100.0)			
悪性高血圧	1,452	5				1,457	7	397	1,861
(%)	(99.7)	(0.3)				(100.0)			
糖尿病性腎症	70,221	120	10		3	70,354	179	18,893	89,426
(%)	(99.8)	(0.2)	(0.0)		(0.0)	(100.0)			
SLE腎炎	1,183	15				1,198	3	303	1,504
(%)	(98.7)	(1.3)				(100.0)			
アミロイド腎	239					239		75	314
(%)	(100.0)					(100.0)			
痛風腎	626	2				628	2	148	778
(%)	(99.7)	(0.3)				(100.0)			
先天性代謝異常による腎不全	132	11				143		30	173
(%)	(92.3)	(7.7)				(100.0)			
腎・尿路結核	111	1				112		29	141
(%)	(99.1)	(0.9)				(100.0)			
腎・尿路結石	344					344		77	421
(%)	(100.0)					(100.0)			
腎・尿路腫瘍	548	4				552		126	678
(%)	(99.3)	(0.7)				(100.0)			
閉塞性尿路障害	400	5				405	1	88	494
(%)	(98.8)	(1.2)				(100.0)			
骨髄腫	149					149	1	40	190
(%)	(100.0)					(100.0)			
腎形成不全	309	26	1			336		64	400
(%)	(92.0)	(7.7)	(0.3)			(100.0)			
不明	16,184	110	7			16,301	83	4,792	21,176
(%)	(99.3)	(0.7)	(0.0)			(100.0)			
再導入	423	743	27	2		1,195	1	160	1,356
(%)	(35.4)	(62.2)	(2.3)	(0.2)		(100.0)			
その他	4,248	67	2			4,317	17	1,381	5,715
(%)	(98.4)	(1.6)	(0.0)			(100.0)			
合計	174,522	1,774	87	3	3	176,389	480	46,976	223,845
(%)	(98.9)	(1.0)	(0.0)	(0.0)	(0.0)	(100.0)			
記載なし	7					7			7
(%)	(100.0)					(100.0)			
総計	174,529	1,774	87	3	3	176,396	480	46,976	223,852
(%)	(98.9)	(1.0)	(0.0)	(0.0)	(0.0)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査