

表 1757 腎移植の回数 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
慢性糸球体腎炎	13,131	285	16	2		13,434	34	3,994	17,462
(%)	(97.7)	(2.1)	(0.1)	(0.0)		(100.0)			
慢性腎盂腎炎	367	3	1			371	1	112	484
(%)	(98.9)	(0.8)	(0.3)			(100.0)			
急速進行性糸球体腎炎	269	3				272		81	353
(%)	(98.9)	(1.1)				(100.0)			
妊娠腎/妊娠中毒症	294	7				301	1	99	401
(%)	(97.7)	(2.3)				(100.0)			
その他の分類不能の腎炎	196	6	1			203	2	56	261
(%)	(96.6)	(3.0)	(0.5)			(100.0)			
多発性嚢胞腎	1,386	7				1,393	2	485	1,880
(%)	(99.5)	(0.5)				(100.0)			
腎硬化症	2,910	8	1			2,919	8	893	3,820
(%)	(99.7)	(0.3)	(0.0)			(100.0)			
悪性高血圧	304	1				305	1	85	391
(%)	(99.7)	(0.3)				(100.0)			
糖尿病性腎症	12,789	27			1	12,817	39	4,104	16,960
(%)	(99.8)	(0.2)			(0.0)	(100.0)			
SLE腎炎	335	3				338	3	85	426
(%)	(99.1)	(0.9)				(100.0)			
アミロイド腎	66					66		8	74
(%)	(100.0)					(100.0)			
痛風腎	125	1				126		40	166
(%)	(99.2)	(0.8)				(100.0)			
先天性代謝異常による腎不全	29	5				34		15	49
(%)	(85.3)	(14.7)				(100.0)			
腎・尿路結核	26					26		5	31
(%)	(100.0)					(100.0)			
腎・尿路結石	64					64		15	79
(%)	(100.0)					(100.0)			
腎・尿路腫瘍	86					86		33	119
(%)	(100.0)					(100.0)			
閉塞性尿路障害	75	2				77		27	104
(%)	(97.4)	(2.6)				(100.0)			
骨髄腫	25					25		8	33
(%)	(100.0)					(100.0)			
腎形成不全	92	7				99		37	136
(%)	(92.9)	(7.1)				(100.0)			
不明	3,459	46	2			3,507	35	1,161	4,703
(%)	(98.6)	(1.3)	(0.1)			(100.0)			
再導入	114	283	18	1		416		67	483
(%)	(27.4)	(68.0)	(4.3)	(0.2)		(100.0)			
その他	805	20	1			826	3	331	1,160
(%)	(97.5)	(2.4)	(0.1)			(100.0)			
合計	36,947	714	40	3	1	37,705	129	11,741	49,575
(%)	(98.0)	(1.9)	(0.1)	(0.0)	(0.0)	(100.0)			
記載なし	6					6			6
(%)	(100.0)					(100.0)			
総計	36,953	714	40	3	1	37,711	129	11,741	49,581
(%)	(98.0)	(1.9)	(0.1)	(0.0)	(0.0)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査