

表 1785 腎移植の回数 原疾患別（腹膜透析患者）

原疾患	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
慢性糸球体腎炎	1,919	8				1,927	3	900	2,830
(%)	(99.6)	(0.4)				(100.0)			
慢性腎盂腎炎	76					76		21	97
(%)	(100.0)					(100.0)			
急速進行性糸球体腎炎	43	1				44		22	66
(%)	(97.7)	(2.3)				(100.0)			
妊娠腎/妊娠中毒症	25					25		2	27
(%)	(100.0)					(100.0)			
その他の分類不能の腎炎	56					56		22	78
(%)	(100.0)					(100.0)			
多発性嚢胞腎	164					164	1	98	263
(%)	(100.0)					(100.0)			
腎硬化症	737	4				741	1	353	1,095
(%)	(99.5)	(0.5)				(100.0)			
悪性高血圧	71					71		36	107
(%)	(100.0)					(100.0)			
糖尿病性腎症	1,845	4				1,849	4	946	2,799
(%)	(99.8)	(0.2)				(100.0)			
SLE腎炎	30					30		16	46
(%)	(100.0)					(100.0)			
アミロイド腎	8					8		7	15
(%)	(100.0)					(100.0)			
痛風腎	25					25		7	32
(%)	(100.0)					(100.0)			
先天性代謝異常による腎不全	11	1				12		5	17
(%)	(91.7)	(8.3)				(100.0)			
腎・尿路結核								2	2
(%)									
腎・尿路結石	13	1				14		7	21
(%)	(92.9)	(7.1)				(100.0)			
腎・尿路腫瘍	11					11		3	14
(%)	(100.0)					(100.0)			
閉塞性尿路障害	13					13		8	21
(%)	(100.0)					(100.0)			
骨髄腫	3					3		4	7
(%)	(100.0)					(100.0)			
腎形成不全	46					46		26	72
(%)	(100.0)					(100.0)			
不明	538	2				540	5	297	842
(%)	(99.6)	(0.4)				(100.0)			
再導入	10	27		1		38		17	55
(%)	(26.3)	(71.1)		(2.6)		(100.0)			
その他	171	1				172		91	263
(%)	(99.4)	(0.6)				(100.0)			
合計	5,815	49		1		5,865	14	2,890	8,769
(%)	(99.1)	(0.8)		(0.0)		(100.0)			
記載なし									
(%)									
総計	5,815	49		1		5,865	14	2,890	8,769
(%)	(99.1)	(0.8)		(0.0)		(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査