

表 1805 週透析回数(回/週) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	1回	2回	3回	4回	5回	6回	7回	合計	記載なし	総計	平均	標準偏差
施設血液透析	561	5,356	223,853	637	16	55	1	230,479	18,246	248,725	2.98	0.19
(%)	(0.2)	(2.3)	(97.1)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
Off-line HDF	8	36	4,836	19		1		4,900	432	5,332	2.99	0.14
(%)	(0.2)	(0.7)	(98.7)	(0.4)		(0.0)		(100.0)				
On-line HDF	44	411	41,006	210	4	4		41,679	2,848	44,527	2.99	0.14
(%)	(0.1)	(1.0)	(98.4)	(0.5)	(0.0)	(0.0)		(100.0)				
Push/Pull HDF			88					88	22	110	3.00	
(%)			(100.0)					(100.0)				
ハイオフィльтраーション		2	249	3				254	13	267	3.00	0.14
(%)		(0.8)	(98.0)	(1.2)				(100.0)				
IHDF	4	65	3,402	10				3,481	59	3,540	2.98	0.16
(%)	(0.1)	(1.9)	(97.7)	(0.3)				(100.0)				
血液濾過	1		12					13	3	16	2.85	0.55
(%)	(7.7)		(92.3)					(100.0)				
血液吸着透析 (リクセル等使用)	1	2	1,325	4				1,332	44	1,376	3.00	0.08
(%)	(0.1)	(0.2)	(99.5)	(0.3)				(100.0)				
在宅血液透析	12	3	128	116	76	79	28	442	113	555	4.38	1.35
(%)	(2.7)	(0.7)	(29.0)	(26.2)	(17.2)	(17.9)	(6.3)	(100.0)				
腹膜透析 (CAPD)	581	69	2					652	4,250	4,902	1.11	0.33
(%)	(89.1)	(10.6)	(0.3)					(100.0)				
腹膜透析 (APD)	364	30	1					395	2,380	2,775	1.08	0.28
(%)	(92.2)	(7.6)	(0.3)					(100.0)				
腹膜透析 (CCPD)	278	15						293	799	1,092	1.05	0.22
(%)	(94.9)	(5.1)						(100.0)				
合計	1,854	5,989	274,902	999	96	139	29	284,008	29,209	313,217	2.97	0.24
(%)	(0.7)	(2.1)	(96.8)	(0.4)	(0.0)	(0.0)	(0.0)	(100.0)				
記載なし												
(%)												
総計	1,854	5,989	274,902	999	96	139	29	284,008	29,209	313,217	2.97	0.24
(%)	(0.7)	(2.1)	(96.8)	(0.4)	(0.0)	(0.0)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査