

表 1806 週透析回数(回/週) 併用の状況別 (透析患者全体)

| 併用の状況              | 1回      | 2回      | 3回      | 4回    | 5回    | 6回    | 7回    | 合計      | 記載なし   | 総計      | 平均   | 標準偏差 |
|--------------------|---------|---------|---------|-------|-------|-------|-------|---------|--------|---------|------|------|
| HD(F)単独            | 626     | 5,832   | 274,688 | 998   | 96    | 137   | 28    | 282,405 | 21,758 | 304,163 | 2.98 | 0.20 |
| (%)                | (0.2)   | (2.1)   | (97.3)  | (0.4) | (0.0) | (0.0) | (0.0) | (100.0) |        |         |      |      |
| PD単独               |         |         |         |       |       |       |       |         | 7,104  | 7,104   |      |      |
| (%)                |         |         |         |       |       |       |       |         |        |         |      |      |
| HD(F)単独.PDカテあり     |         | 2       | 165     | 1     |       | 1     |       | 169     | 13     | 182     | 3.01 | 0.27 |
| (%)                |         | (1.2)   | (97.6)  | (0.6) |       | (0.6) |       | (100.0) |        |         |      |      |
| PD.HD併用週1<br>HD(F) | 1,202   |         |         |       |       |       |       | 1,202   | 268    | 1,470   | 1.00 |      |
| (%)                | (100.0) |         |         |       |       |       |       | (100.0) |        |         |      |      |
| PD.HD併用週2<br>HD(F) |         | 153     |         |       |       |       |       | 153     | 19     | 172     | 2.00 |      |
| (%)                |         | (100.0) |         |       |       |       |       | (100.0) |        |         |      |      |
| PD.HD併用週3<br>HD(F) |         |         | 26      |       |       |       |       | 26      | 3      | 29      | 3.00 |      |
| (%)                |         |         | (100.0) |       |       |       |       | (100.0) |        |         |      |      |
| PD.HD併用週4<br>HD(F) |         |         |         |       |       |       |       |         |        |         |      |      |
| (%)                |         |         |         |       |       |       |       |         |        |         |      |      |
| PD.HD併用その他         | 26      | 2       | 5       |       |       |       |       | 33      | 38     | 71      | 1.36 | 0.74 |
| (%)                | (78.8)  | (6.1)   | (15.2)  |       |       |       |       | (100.0) |        |         |      |      |
| 合計                 | 1,854   | 5,989   | 274,884 | 999   | 96    | 138   | 28    | 283,988 | 29,203 | 313,191 | 2.97 | 0.24 |
| (%)                | (0.7)   | (2.1)   | (96.8)  | (0.4) | (0.0) | (0.0) | (0.0) | (100.0) |        |         |      |      |
| 不明                 |         |         | 18      |       |       | 1     | 1     | 20      | 5      | 25      | 3.35 | 1.09 |
| (%)                |         |         | (90.0)  |       |       | (5.0) | (5.0) | (100.0) |        |         |      |      |
| 記載なし               |         |         |         |       |       |       |       |         | 1      | 1       |      |      |
| (%)                |         |         |         |       |       |       |       |         |        |         |      |      |
| 総計                 | 1,854   | 5,989   | 274,902 | 999   | 96    | 139   | 29    | 284,008 | 29,209 | 313,217 | 2.97 | 0.24 |
| (%)                | (0.7)   | (2.1)   | (96.8)  | (0.4) | (0.0) | (0.0) | (0.0) | (100.0) |        |         |      |      |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査