

表 1825 透析時間(時間/回) 原疾患別(施設血液透析患者 透析回数週3回)

原疾患	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	101	5,000	4,062	43,635	4,023	5,355	260	398	17	201	63,052	422	63,474	4.04	0.53
(%)	(0.2)	(7.9)	(6.4)	(69.2)	(6.4)	(8.5)	(0.4)	(0.6)	(0.0)	(0.3)	(100.0)				
慢性腎盂腎炎	4	221	158	1,360	99	154	4	20		4	2,024	16	2,040	3.99	0.55
(%)	(0.2)	(10.9)	(7.8)	(67.2)	(4.9)	(7.6)	(0.2)	(1.0)		(0.2)	(100.0)				
急速進行性糸球体腎炎	5	235	129	1,212	61	106	2	11	1	2	1,764	7	1,771	3.93	0.52
(%)	(0.3)	(13.3)	(7.3)	(68.7)	(3.5)	(6.0)	(0.1)	(0.6)	(0.1)	(0.1)	(100.0)				
妊娠腎/妊娠中毒症	2	48	69	712	73	59	5	6		4	978	3	981	4.06	0.50
(%)	(0.2)	(4.9)	(7.1)	(72.8)	(7.5)	(6.0)	(0.5)	(0.6)		(0.4)	(100.0)				
その他の分類不能の腎炎	2	102	58	606	64	75	3	1	1	12	924	5	929	4.04	0.67
(%)	(0.2)	(11.0)	(6.3)	(65.6)	(6.9)	(8.1)	(0.3)	(0.1)	(0.1)	(1.3)	(100.0)				
多発性嚢胞腎	8	761	657	5,513	418	551	19	29		27	7,983	33	8,016	3.99	0.52
(%)	(0.1)	(9.5)	(8.2)	(69.1)	(5.2)	(6.9)	(0.2)	(0.4)		(0.3)	(100.0)				
腎硬化症	66	3,160	2,111	14,462	826	1,100	27	95	4	43	21,894	113	22,007	3.90	0.52
(%)	(0.3)	(14.4)	(9.6)	(66.1)	(3.8)	(5.0)	(0.1)	(0.4)	(0.0)	(0.2)	(100.0)				
悪性高血圧	1	199	157	1,214	109	157	4	6		4	1,851	10	1,861	3.99	0.52
(%)	(0.1)	(10.8)	(8.5)	(65.6)	(5.9)	(8.5)	(0.2)	(0.3)		(0.2)	(100.0)				
糖尿病性腎症	164	9,487	6,929	61,601	4,077	5,838	184	423	15	163	88,881	545	89,426	3.97	0.51
(%)	(0.2)	(10.7)	(7.8)	(69.3)	(4.6)	(6.6)	(0.2)	(0.5)	(0.0)	(0.2)	(100.0)				
SLE腎炎	3	166	108	1,020	87	97	3	5	1	3	1,493	11	1,504	3.97	0.52
(%)	(0.2)	(11.1)	(7.2)	(68.3)	(5.8)	(6.5)	(0.2)	(0.3)	(0.1)	(0.2)	(100.0)				
アミロイド腎	1	46	31	198	14	19		1			310	4	314	3.90	0.50
(%)	(0.3)	(14.8)	(10.0)	(63.9)	(4.5)	(6.1)		(0.3)			(100.0)				
痛風腎	1	63	53	556	38	49	1	4		5	770	8	778	4.02	0.53
(%)	(0.1)	(8.2)	(6.9)	(72.2)	(4.9)	(6.4)	(0.1)	(0.5)		(0.6)	(100.0)				
先天性代謝異常による腎不全	1	15	10	114	12	19		1			172	1	173	4.04	0.52
(%)	(0.6)	(8.7)	(5.8)	(66.3)	(7.0)	(11.0)		(0.6)			(100.0)				
腎・尿路結核	1	22	14	90	4	9					140	1	141	3.87	0.50
(%)	(0.7)	(15.7)	(10.0)	(64.3)	(2.9)	(6.4)					(100.0)				
腎・尿路結石	1	56	39	283	16	20	1	1		2	419	2	421	3.92	0.55
(%)	(0.2)	(13.4)	(9.3)	(67.5)	(3.8)	(4.8)	(0.2)	(0.2)		(0.5)	(100.0)				
腎・尿路腫瘍	3	85	53	476	16	34		4		2	673	5	678	3.92	0.52
(%)	(0.4)	(12.6)	(7.9)	(70.7)	(2.4)	(5.1)		(0.6)		(0.3)	(100.0)				
閉塞性尿路障害	3	64	38	321	27	36	1			3	493	1	494	3.96	0.61
(%)	(0.6)	(13.0)	(7.7)	(65.1)	(5.5)	(7.3)	(0.2)			(0.6)	(100.0)				
骨髄腫		58	17	101	4	9		1			190		190	3.73	0.57
(%)		(30.5)	(8.9)	(53.2)	(2.1)	(4.7)		(0.5)			(100.0)				
腎形成不全		24	30	277	33	31		2		2	399	1	400	4.06	0.51
(%)		(6.0)	(7.5)	(69.4)	(8.3)	(7.8)		(0.5)		(0.5)	(100.0)				
不明	70	3,143	1,990	13,930	863	900	45	55	3	44	21,043	133	21,176	3.89	0.52
(%)	(0.3)	(14.9)	(9.5)	(66.2)	(4.1)	(4.3)	(0.2)	(0.3)	(0.0)	(0.2)	(100.0)				
再導入	2	99	85	907	109	125	4	11	1	6	1,349	7	1,356	4.08	0.56
(%)	(0.1)	(7.3)	(6.3)	(67.2)	(8.1)	(9.3)	(0.3)	(0.8)	(0.1)	(0.4)	(100.0)				
その他	22	742	425	3,822	281	325	16	26	1	23	5,683	32	5,715	3.95	0.55
(%)	(0.4)	(13.1)	(7.5)	(67.3)	(4.9)	(5.7)	(0.3)	(0.5)	(0.0)	(0.4)	(100.0)				
合計	461	23,796	17,223	152,410	11,254	15,068	579	1,100	44	550	222,485	1,360	223,845	3.98	0.52
(%)	(0.2)	(10.7)	(7.7)	(68.5)	(5.1)	(6.8)	(0.3)	(0.5)	(0.0)	(0.2)	(100.0)				
記載なし		3		4							7		7	3.57	0.53
(%)		(42.9)		(57.1)							(100.0)				
総計	461	23,799	17,223	152,414	11,254	15,068	579	1,100	44	550	222,492	1,360	223,852	3.98	0.52
(%)	(0.2)	(10.7)	(7.7)	(68.5)	(5.1)	(6.8)	(0.3)	(0.5)	(0.0)	(0.2)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査