

表 1876 血流量(ml/分) 性別（施設血液透析患者 透析回数週3回）

性別	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	280～	300～	350～	400～	合計	記載なし	総計	平均	標準偏差
男性	22	362	972	7,867	2,303	14,417	61,476	22,439	22,868	2,125	2,360	3,947	399	378	141,935	2,505	144,440	211.75	36.46
(%)	(0.0)	(0.3)	(0.7)	(5.5)	(1.6)	(10.2)	(43.3)	(15.8)	(16.1)	(1.5)	(1.7)	(2.8)	(0.3)	(0.3)	(100.0)				
女性	26	347	1,119	9,523	2,996	14,134	35,996	7,018	5,098	360	373	601	68	196	77,855	1,557	79,412	194.94	33.34
(%)	(0.0)	(0.4)	(1.4)	(12.2)	(3.8)	(18.2)	(46.2)	(9.0)	(6.5)	(0.5)	(0.5)	(0.8)	(0.1)	(0.3)	(100.0)				
合計	48	709	2,091	17,390	5,299	28,551	97,472	29,457	27,966	2,485	2,733	4,548	467	574	219,790	4,062	223,852	205.79	36.29
(%)	(0.0)	(0.3)	(1.0)	(7.9)	(2.4)	(13.0)	(44.3)	(13.4)	(12.7)	(1.1)	(1.2)	(2.1)	(0.2)	(0.3)	(100.0)				
記載なし																			
(%)																			
総計	48	709	2,091	17,390	5,299	28,551	97,472	29,457	27,966	2,485	2,733	4,548	467	574	219,790	4,062	223,852	205.79	36.29
(%)	(0.0)	(0.3)	(1.0)	(7.9)	(2.4)	(13.0)	(44.3)	(13.4)	(12.7)	(1.1)	(1.2)	(2.1)	(0.2)	(0.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査