

表 1931 HDF希釈方法 原疾患別（オフラインHDF 透析回数週3回）

原疾患	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
慢性糸球体腎炎	225	1,533		4	1,762		236	1,998
(%)	(12.8)	(87.0)		(0.2)	(100.0)			
慢性腎盂腎炎	3	41			44		10	54
(%)	(6.8)	(93.2)			(100.0)			
急速進行性糸球体腎炎	5	25			30		1	31
(%)	(16.7)	(83.3)			(100.0)			
妊娠腎/妊娠中毒症	2	43			45		10	55
(%)	(4.4)	(95.6)			(100.0)			
その他の分類不能の腎炎	4	15			19		4	23
(%)	(21.1)	(78.9)			(100.0)			
多発性嚢胞腎	14	119			133		18	151
(%)	(10.5)	(89.5)			(100.0)			
腎硬化症	16	192			208		38	246
(%)	(7.7)	(92.3)			(100.0)			
悪性高血圧	3	23			26		8	34
(%)	(11.5)	(88.5)			(100.0)			
糖尿病性腎症	155	1,135		2	1,292		198	1,490
(%)	(12.0)	(87.8)		(0.2)	(100.0)			
SLE腎炎	3	40			43		11	54
(%)	(7.0)	(93.0)			(100.0)			
アミロイド腎		10			10		2	12
(%)		(100.0)			(100.0)			
痛風腎	1	15			16		3	19
(%)	(6.3)	(93.8)			(100.0)			
先天性代謝異常による腎不全	1	3			4			4
(%)	(25.0)	(75.0)			(100.0)			
腎・尿路結核		1			1			1
(%)		(100.0)			(100.0)			
腎・尿路結石		7			7		1	8
(%)		(100.0)			(100.0)			
腎・尿路腫瘍	3	10			13		5	18
(%)	(23.1)	(76.9)			(100.0)			
閉塞性尿路障害	2	6			8		1	9
(%)	(25.0)	(75.0)			(100.0)			
骨髄腫	2	5			7		2	9
(%)	(28.6)	(71.4)			(100.0)			
腎形成不全	1	13			14		1	15
(%)	(7.1)	(92.9)			(100.0)			
不明	28	329			357	1	72	430
(%)	(7.8)	(92.2)			(100.0)			
再導入	5	54			59		6	65
(%)	(8.5)	(91.5)			(100.0)			
その他	11	82			93		17	110
(%)	(11.8)	(88.2)			(100.0)			
合計	484	3,701		6	4,191	1	644	4,836
(%)	(11.5)	(88.3)		(0.1)	(100.0)			
記載なし								
(%)								
総計	484	3,701		6	4,191	1	644	4,836
(%)	(11.5)	(88.3)		(0.1)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査