

表 1939 HDF希釈方法 透析時間(時間/回)別 (オフラインHDF 透析回数週3回)

透析時間	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
3.0未満	1	3			4			4
(%)	(25.0)	(75.0)			(100.0)			
3.0～	40	190			230		51	281
(%)	(17.4)	(82.6)			(100.0)			
3.5～	12	194			206		30	236
(%)	(5.8)	(94.2)			(100.0)			
4.0～	344	2,507		3	2,854	1	439	3,294
(%)	(12.1)	(87.8)		(0.1)	(100.0)			
4.5～	40	349		2	391		40	431
(%)	(10.2)	(89.3)		(0.5)	(100.0)			
5.0～	40	394		1	435		59	494
(%)	(9.2)	(90.6)		(0.2)	(100.0)			
5.5～	2	15			17		3	20
(%)	(11.8)	(88.2)			(100.0)			
6.0～	3	32			35		3	38
(%)	(8.6)	(91.4)			(100.0)			
6.5～		6			6			6
(%)		(100.0)			(100.0)			
7.0～	1				1		2	3
(%)	(100.0)				(100.0)			
合計	483	3,690		6	4,179	1	627	4,807
(%)	(11.6)	(88.3)		(0.1)	(100.0)			
記載なし	1	11			12		17	29
(%)	(8.3)	(91.7)			(100.0)			
総計	484	3,701		6	4,191	1	644	4,836
(%)	(11.5)	(88.3)		(0.1)	(100.0)			
平均	4.05	4.12		4.36	4.11	4.00	4.06	4.10
標準偏差	0.50	0.49		0.39	0.49		0.53	0.50

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査