

表 1952 HDF希釈方法 透析前CRP濃度(mg/dL)別（オフラインHDF 透析回数週3回）

透析前CRP濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
0.2未満	171	1,578		2	1,751	1	228	1,980
(%)	(9.8)	(90.1)		(0.1)	(100.0)			
0.2～	100	660		1	761		86	847
(%)	(13.1)	(86.7)		(0.1)	(100.0)			
0.5～	45	420		2	467		46	513
(%)	(9.6)	(89.9)		(0.4)	(100.0)			
1.0～	30	239			269		58	327
(%)	(11.2)	(88.8)			(100.0)			
2.0～	20	127			147		31	178
(%)	(13.6)	(86.4)			(100.0)			
4.0～	10	54			64		5	69
(%)	(15.6)	(84.4)			(100.0)			
6.0～	6	31			37		6	43
(%)	(16.2)	(83.8)			(100.0)			
8.0～		15			15		3	18
(%)		(100.0)			(100.0)			
10.0～	3	12			15		3	18
(%)	(20.0)	(80.0)			(100.0)			
15.0～	1	8			9		1	10
(%)	(11.1)	(88.9)			(100.0)			
合計	386	3,144		5	3,535	1	467	4,003
(%)	(10.9)	(88.9)		(0.1)	(100.0)			
記載なし	98	557		1	656		177	833
(%)	(14.9)	(84.9)		(0.2)	(100.0)			
総計	484	3,701		6	4,191	1	644	4,836
(%)	(11.5)	(88.3)		(0.1)	(100.0)			
平均	0.80	0.72		0.36	0.73	0.09	0.87	0.74
標準偏差	1.76	1.74		0.37	1.74		2.25	1.81

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査