

表 1953 HDF希釈方法 透析前補正カルシウム濃度(mg/dL)別（オフラインHDF 透析回数週3回）

透析前補正 カルシウム濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
7.0未満	2	17			19		5	24
(%)	(10.5)	(89.5)			(100.0)			
7.0～	5	18			23		5	28
(%)	(21.7)	(78.3)			(100.0)			
7.5～	9	85			94		12	106
(%)	(9.6)	(90.4)			(100.0)			
8.0～	27	215		1	243		26	269
(%)	(11.1)	(88.5)		(0.4)	(100.0)			
8.4～	8	67			75		10	85
(%)	(10.7)	(89.3)			(100.0)			
8.5～	92	766			858		117	975
(%)	(10.7)	(89.3)			(100.0)			
9.0～	149	1,025		2	1,176	1	185	1,362
(%)	(12.7)	(87.2)		(0.2)	(100.0)			
9.5～	101	777		2	880		121	1,001
(%)	(11.5)	(88.3)		(0.2)	(100.0)			
10.0～	13	105			118		11	129
(%)	(11.0)	(89.0)			(100.0)			
10.1～	29	271			300		39	339
(%)	(9.7)	(90.3)			(100.0)			
10.5～	26	148		1	175		29	204
(%)	(14.9)	(84.6)		(0.6)	(100.0)			
11.0～	5	60			65		13	78
(%)	(7.7)	(92.3)			(100.0)			
11.5～	2	19			21		4	25
(%)	(9.5)	(90.5)			(100.0)			
12.0～	1	12			13			13
(%)	(7.7)	(92.3)			(100.0)			
合計	469	3,585		6	4,060	1	577	4,638
(%)	(11.6)	(88.3)		(0.1)	(100.0)			
記載なし	15	116			131		67	198
(%)	(11.5)	(88.5)			(100.0)			
総計	484	3,701		6	4,191	1	644	4,836
(%)	(11.5)	(88.3)		(0.1)	(100.0)			
平均	9.27	9.29		9.35	9.29	9.10	9.30	9.29
標準偏差	0.80	0.80		0.76	0.80		0.79	0.80

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査