

表 1954 HDF希釈方法 透析前リン濃度(mg/dL)別（オフラインHDF 透析回数週3回）

透析前リン濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
3未満	20	155			175		32	207
(%)	(11.4)	(88.6)			(100.0)			
3.0～	33	160			193		32	225
(%)	(17.1)	(82.9)			(100.0)			
3.5～	32	282		1	315	1	37	353
(%)	(10.2)	(89.5)		(0.3)	(100.0)			
4.0～	58	420		1	479		82	561
(%)	(12.1)	(87.7)		(0.2)	(100.0)			
4.5～	58	538		1	597		90	687
(%)	(9.7)	(90.1)		(0.2)	(100.0)			
5.0～	64	543		1	608		85	693
(%)	(10.5)	(89.3)		(0.2)	(100.0)			
5.5～	61	463			524		73	597
(%)	(11.6)	(88.4)			(100.0)			
6.0～	8	71		1	80		12	92
(%)	(10.0)	(88.8)		(1.3)	(100.0)			
6.1～	26	278			304		31	335
(%)	(8.6)	(91.4)			(100.0)			
6.5～	48	266			314		43	357
(%)	(15.3)	(84.7)			(100.0)			
7.0～	23	159			182		23	205
(%)	(12.6)	(87.4)			(100.0)			
7.5～	11	90			101		16	117
(%)	(10.9)	(89.1)			(100.0)			
8.0～	6	72			78		14	92
(%)	(7.7)	(92.3)			(100.0)			
8.5～	7	32			39		6	45
(%)	(17.9)	(82.1)			(100.0)			
9.0～	13	66		1	80		16	96
(%)	(16.3)	(82.5)		(1.3)	(100.0)			
合計	468	3,595		6	4,069	1	592	4,662
(%)	(11.5)	(88.4)		(0.1)	(100.0)			
記載なし	16	106			122		52	174
(%)	(13.1)	(86.9)			(100.0)			
総計	484	3,701		6	4,191	1	644	4,836
(%)	(11.5)	(88.3)		(0.1)	(100.0)			
平均	5.33	5.29		5.67	5.29	3.50	5.25	5.29
標準偏差	1.60	1.49		2.37	1.51		1.58	1.52

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査