

表 1956 HDF希釈方法 透析前ヘモグロビン濃度(g/dL)別（オフラインHDF 透析回数週3回）

透析前ヘモグロビン濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
5未満	1	2			3		1	4
(%)	(33.3)	(66.7)			(100.0)			
5～		7			7			7
(%)		(100.0)			(100.0)			
6～	2	16			18		1	19
(%)	(11.1)	(88.9)			(100.0)			
7～	8	60			68		8	76
(%)	(11.8)	(88.2)			(100.0)			
8～	29	183			212		47	259
(%)	(13.7)	(86.3)			(100.0)			
9～	78	594		3	675		102	777
(%)	(11.6)	(88.0)		(0.4)	(100.0)			
10～	165	1,137		1	1,303	1	185	1,489
(%)	(12.7)	(87.3)		(0.1)	(100.0)			
11～	102	1,018		2	1,122		149	1,271
(%)	(9.1)	(90.7)		(0.2)	(100.0)			
12～	56	406			462		70	532
(%)	(12.1)	(87.9)			(100.0)			
13～	14	120			134		17	151
(%)	(10.4)	(89.6)			(100.0)			
14～	11	35			46		5	51
(%)	(23.9)	(76.1)			(100.0)			
合計	466	3,578		6	4,050	1	585	4,636
(%)	(11.5)	(88.3)		(0.1)	(100.0)			
記載なし	18	123			141		59	200
(%)	(12.8)	(87.2)			(100.0)			
総計	484	3,701		6	4,191	1	644	4,836
(%)	(11.5)	(88.3)		(0.1)	(100.0)			
平均	10.73	10.75		10.42	10.74	10.10	10.68	10.74
標準偏差	1.38	1.31		1.04	1.32		1.32	1.32

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査