

表 1966 HDF希釈方法 原疾患別（オンラインHDF 透析回数週3回）

原疾患	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
慢性糸球体腎炎	12,484	647	1	61	13,193		1,158	14,351
(%)	(94.6)	(4.9)	(0.0)	(0.5)	(100.0)			
慢性腎盂腎炎	358	13		3	374		28	402
(%)	(95.7)	(3.5)		(0.8)	(100.0)			
急速進行性糸球体腎炎	259	8		1	268		29	297
(%)	(96.6)	(3.0)		(0.4)	(100.0)			
妊娠腎/妊娠中毒症	283	16			299		27	326
(%)	(94.6)	(5.4)			(100.0)			
その他の分類不能の腎炎	171	15		2	188		21	209
(%)	(91.0)	(8.0)		(1.1)	(100.0)			
多発性嚢胞腎	1,431	69		8	1,508		114	1,622
(%)	(94.9)	(4.6)		(0.5)	(100.0)			
腎硬化症	2,833	113		28	2,974		224	3,198
(%)	(95.3)	(3.8)		(0.9)	(100.0)			
悪性高血圧	302	12		2	316		16	332
(%)	(95.6)	(3.8)		(0.6)	(100.0)			
糖尿病性腎症	12,265	521		135	12,921		999	13,920
(%)	(94.9)	(4.0)		(1.0)	(100.0)			
SLE腎炎	313	15			328		25	353
(%)	(95.4)	(4.6)			(100.0)			
アミロイド腎	51	2			53		4	57
(%)	(96.2)	(3.8)			(100.0)			
痛風腎	120	8		1	129		8	137
(%)	(93.0)	(6.2)		(0.8)	(100.0)			
先天性代謝異常による腎不全	36	4			40		3	43
(%)	(90.0)	(10.0)			(100.0)			
腎・尿路結核	24	1			25		1	26
(%)	(96.0)	(4.0)			(100.0)			
腎・尿路結石	61	1		1	63		4	67
(%)	(96.8)	(1.6)		(1.6)	(100.0)			
腎・尿路腫瘍	81	2			83		7	90
(%)	(97.6)	(2.4)			(100.0)			
閉塞性尿路障害	76	2			78		9	87
(%)	(97.4)	(2.6)			(100.0)			
骨髄腫	23				23		1	24
(%)	(100.0)				(100.0)			
腎形成不全	96	2			98		10	108
(%)	(98.0)	(2.0)			(100.0)			
不明	3,536	115		56	3,707		278	3,985
(%)	(95.4)	(3.1)		(1.5)	(100.0)			
再導入	352	19		4	375		22	397
(%)	(93.9)	(5.1)		(1.1)	(100.0)			
その他	833	32		8	873		96	969
(%)	(95.4)	(3.7)		(0.9)	(100.0)			
合計	35,988	1,617	1	310	37,916		3,084	41,000
(%)	(94.9)	(4.3)	(0.0)	(0.8)	(100.0)			
記載なし	6				6			6
(%)	(100.0)				(100.0)			
総計	35,994	1,617	1	310	37,922		3,084	41,006
(%)	(94.9)	(4.3)	(0.0)	(0.8)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査