

表 1974 HDF希釈方法 透析時間(時間/回)別 (オンラインHDF 透析回数週3回)

透析時間	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
3.0未満	53	1			54		2	56
(%)	(98.1)	(1.9)			(100.0)			
3.0～	2,075	124		23	2,222		121	2,343
(%)	(93.4)	(5.6)		(1.0)	(100.0)			
3.5～	2,274	103		19	2,396		145	2,541
(%)	(94.9)	(4.3)		(0.8)	(100.0)			
4.0～	24,117	1,044	1	236	25,398		2,087	27,485
(%)	(95.0)	(4.1)	(0.0)	(0.9)	(100.0)			
4.5～	3,311	164		20	3,495		263	3,758
(%)	(94.7)	(4.7)		(0.6)	(100.0)			
5.0～	3,655	163		12	3,830		269	4,099
(%)	(95.4)	(4.3)		(0.3)	(100.0)			
5.5～	167	6			173		9	182
(%)	(96.5)	(3.5)			(100.0)			
6.0～	245	9			254		18	272
(%)	(96.5)	(3.5)			(100.0)			
6.5～	11				11			11
(%)	(100.0)				(100.0)			
7.0～	29				29		1	30
(%)	(100.0)				(100.0)			
合計	35,937	1,614	1	310	37,862		2,915	40,777
(%)	(94.9)	(4.3)	(0.0)	(0.8)	(100.0)			
記載なし	57	3			60		169	229
(%)	(95.0)	(5.0)			(100.0)			
総計	35,994	1,617	1	310	37,922		3,084	41,006
(%)	(94.9)	(4.3)	(0.0)	(0.8)	(100.0)			
平均	4.10	4.07	4.00	3.97	4.09		4.12	4.10
標準偏差	0.50	0.50		0.37	0.50		0.45	0.49

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査