

表 1988 HDF希釈方法 透析前補正カルシウム濃度(mg/dL)別（オンラインHDF 透析回数週3回）

透析前補正 カルシウム濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
7.0未満	137	2			139		6	145
(%)	(98.6)	(1.4)			(100.0)			
7.0～	207	8		1	216		9	225
(%)	(95.8)	(3.7)		(0.5)	(100.0)			
7.5～	863	26		4	893		79	972
(%)	(96.6)	(2.9)		(0.4)	(100.0)			
8.0～	2,661	100		18	2,779		215	2,994
(%)	(95.8)	(3.6)		(0.6)	(100.0)			
8.4～	698	27	1	4	730		56	786
(%)	(95.6)	(3.7)	(0.1)	(0.5)	(100.0)			
8.5～	8,478	349		72	8,899		730	9,629
(%)	(95.3)	(3.9)		(0.8)	(100.0)			
9.0～	10,626	467		62	11,155		831	11,986
(%)	(95.3)	(4.2)		(0.6)	(100.0)			
9.5～	7,109	331		35	7,475		571	8,046
(%)	(95.1)	(4.4)		(0.5)	(100.0)			
10.0～	832	44		4	880		72	952
(%)	(94.5)	(5.0)		(0.5)	(100.0)			
10.1～	2,127	131		9	2,267		163	2,430
(%)	(93.8)	(5.8)		(0.4)	(100.0)			
10.5～	979	59		3	1,041		84	1,125
(%)	(94.0)	(5.7)		(0.3)	(100.0)			
11.0～	300	21			321		28	349
(%)	(93.5)	(6.5)			(100.0)			
11.5～	107	7			114		4	118
(%)	(93.9)	(6.1)			(100.0)			
12.0～	94				94		2	96
(%)	(100.0)				(100.0)			
合計	35,218	1,572	1	212	37,003		2,850	39,853
(%)	(95.2)	(4.2)	(0.0)	(0.6)	(100.0)			
記載なし	776	45		98	919		234	1,153
(%)	(84.4)	(4.9)		(10.7)	(100.0)			
総計	35,994	1,617	1	310	37,922		3,084	41,006
(%)	(94.9)	(4.3)	(0.0)	(0.8)	(100.0)			
平均	9.18	9.28	8.40	9.06	9.19		9.18	9.19
標準偏差	0.73	0.71		0.58	0.73		0.70	0.73

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査