

表 1989 HDF希釈方法 透析前リン濃度(mg/dL)別 (オンラインHDF 透析回数週3回)

透析前リン濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
3未満	913	48		10	971		85	1,056
(%)	(94.0)	(4.9)		(1.0)	(100.0)			
3.0～	1,284	50		9	1,343		95	1,438
(%)	(95.6)	(3.7)		(0.7)	(100.0)			
3.5～	2,447	106		14	2,567		187	2,754
(%)	(95.3)	(4.1)		(0.5)	(100.0)			
4.0～	3,948	180		19	4,147		319	4,466
(%)	(95.2)	(4.3)		(0.5)	(100.0)			
4.5～	5,068	242		24	5,334		385	5,719
(%)	(95.0)	(4.5)		(0.4)	(100.0)			
5.0～	5,511	249		33	5,793		451	6,244
(%)	(95.1)	(4.3)		(0.6)	(100.0)			
5.5～	5,011	209		27	5,247		378	5,625
(%)	(95.5)	(4.0)		(0.5)	(100.0)			
6.0～	918	46		6	970		75	1,045
(%)	(94.6)	(4.7)		(0.6)	(100.0)			
6.1～	3,063	115	1	25	3,204		259	3,463
(%)	(95.6)	(3.6)	(0.0)	(0.8)	(100.0)			
6.5～	2,699	123		13	2,835		237	3,072
(%)	(95.2)	(4.3)		(0.5)	(100.0)			
7.0～	1,713	77		11	1,801		165	1,966
(%)	(95.1)	(4.3)		(0.6)	(100.0)			
7.5～	1,088	56		6	1,150		87	1,237
(%)	(94.6)	(4.9)		(0.5)	(100.0)			
8.0～	647	21		6	674		57	731
(%)	(96.0)	(3.1)		(0.9)	(100.0)			
8.5～	384	23		4	411		36	447
(%)	(93.4)	(5.6)		(1.0)	(100.0)			
9.0～	637	29		5	671		44	715
(%)	(94.9)	(4.3)		(0.7)	(100.0)			
合計	35,331	1,574	1	212	37,118		2,860	39,978
(%)	(95.2)	(4.2)	(0.0)	(0.6)	(100.0)			
記載なし	663	43		98	804		224	1,028
(%)	(82.5)	(5.3)		(12.2)	(100.0)			
総計	35,994	1,617	1	310	37,922		3,084	41,006
(%)	(94.9)	(4.3)	(0.0)	(0.8)	(100.0)			
平均	5.42	5.41	6.30	5.47	5.42		5.45	5.42
標準偏差	1.42	1.42		1.54	1.42		1.44	1.42

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査