

表 1991 HDF希釈方法 透析前ヘモグロビン濃度(g/dL)別（オンラインHDF 透析回数週3回）

| 透析前ヘモグロビン<br>濃度 | 前希釈    | 後希釈   | 前・後希釈 | その他    | 合計      | 不明 | 記載なし  | 総計     |
|-----------------|--------|-------|-------|--------|---------|----|-------|--------|
| 5未満             | 24     | 1     |       |        | 25      |    | 2     | 27     |
| (%)             | (96.0) | (4.0) |       |        | (100.0) |    |       |        |
| 5～              | 29     | 1     |       |        | 30      |    | 1     | 31     |
| (%)             | (96.7) | (3.3) |       |        | (100.0) |    |       |        |
| 6～              | 81     | 3     |       | 2      | 86      |    | 4     | 90     |
| (%)             | (94.2) | (3.5) |       | (2.3)  | (100.0) |    |       |        |
| 7～              | 346    | 10    |       | 3      | 359     |    | 22    | 381    |
| (%)             | (96.4) | (2.8) |       | (0.8)  | (100.0) |    |       |        |
| 8～              | 1,179  | 69    |       | 9      | 1,257   |    | 118   | 1,375  |
| (%)             | (93.8) | (5.5) |       | (0.7)  | (100.0) |    |       |        |
| 9～              | 4,749  | 232   |       | 31     | 5,012   |    | 379   | 5,391  |
| (%)             | (94.8) | (4.6) |       | (0.6)  | (100.0) |    |       |        |
| 10～             | 11,793 | 526   |       | 87     | 12,406  |    | 1,031 | 13,437 |
| (%)             | (95.1) | (4.2) |       | (0.7)  | (100.0) |    |       |        |
| 11～             | 11,049 | 483   | 1     | 52     | 11,585  |    | 863   | 12,448 |
| (%)             | (95.4) | (4.2) | (0.0) | (0.4)  | (100.0) |    |       |        |
| 12～             | 4,248  | 192   |       | 25     | 4,465   |    | 296   | 4,761  |
| (%)             | (95.1) | (4.3) |       | (0.6)  | (100.0) |    |       |        |
| 13～             | 1,170  | 41    |       | 2      | 1,213   |    | 92    | 1,305  |
| (%)             | (96.5) | (3.4) |       | (0.2)  | (100.0) |    |       |        |
| 14～             | 516    | 15    |       | 1      | 532     |    | 34    | 566    |
| (%)             | (97.0) | (2.8) |       | (0.2)  | (100.0) |    |       |        |
| 合計              | 35,184 | 1,573 | 1     | 212    | 36,970  |    | 2,842 | 39,812 |
| (%)             | (95.2) | (4.3) | (0.0) | (0.6)  | (100.0) |    |       |        |
| 記載なし            | 810    | 44    |       | 98     | 952     |    | 242   | 1,194  |
| (%)             | (85.1) | (4.6) |       | (10.3) | (100.0) |    |       |        |
| 総計              | 35,994 | 1,617 | 1     | 310    | 37,922  |    | 3,084 | 41,006 |
| (%)             | (94.9) | (4.3) | (0.0) | (0.8)  | (100.0) |    |       |        |
| 平均              | 10.92  | 10.84 | 11.10 | 10.68  | 10.92   |    | 10.85 | 10.91  |
| 標準偏差            | 1.24   | 1.19  |       | 1.20   | 1.24    |    | 1.19  | 1.23   |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査