

表 2087 1セッションあたりの置換液量(L) 原疾患別 (オンラインHDF 透析回数週3回 後希釈)

原疾患	1未満	1～	5～	10～	15～	20～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎		49	212	284	58	34	637	10	647	10.15	4.66
(%)		(7.7)	(33.3)	(44.6)	(9.1)	(5.3)	(100.0)				
慢性腎盂腎炎		1	5	5	1	1	13		13	10.15	4.43
(%)		(7.7)	(38.5)	(38.5)	(7.7)	(7.7)	(100.0)				
急速進行性糸球体腎炎			4	4			8		8	9.00	2.56
(%)			(50.0)	(50.0)			(100.0)				
妊娠腎/妊娠中毒症		2	6	7			15	1	16	8.40	3.25
(%)		(13.3)	(40.0)	(46.7)			(100.0)				
その他の分類不能の腎炎		1	6	7		1	15		15	9.53	3.85
(%)		(6.7)	(40.0)	(46.7)		(6.7)	(100.0)				
多発性嚢胞腎		3	17	36	4	9	69		69	11.17	4.63
(%)		(4.3)	(24.6)	(52.2)	(5.8)	(13.0)	(100.0)				
腎硬化症		6	51	35	6	13	111	2	113	10.32	5.26
(%)		(5.4)	(45.9)	(31.5)	(5.4)	(11.7)	(100.0)				
悪性高血圧		1	4	4	1	2	12		12	11.25	5.61
(%)		(8.3)	(33.3)	(33.3)	(8.3)	(16.7)	(100.0)				
糖尿病性腎症		60	229	161	20	49	519	2	521	9.43	4.98
(%)		(11.6)	(44.1)	(31.0)	(3.9)	(9.4)	(100.0)				
SLE腎炎		1	4	7		2	14	1	15	10.79	4.84
(%)		(7.1)	(28.6)	(50.0)		(14.3)	(100.0)				
アミロイド腎			2				2		2	8.00	
(%)			(100.0)				(100.0)				
痛風腎			4	3	1		8		8	10.13	4.19
(%)			(50.0)	(37.5)	(12.5)		(100.0)				
先天性代謝異常による腎不全			2	2			4		4	10.50	2.38
(%)			(50.0)	(50.0)			(100.0)				
腎・尿路結核			1				1		1	7.00	
(%)			(100.0)				(100.0)				
腎・尿路結石				1			1		1	10.00	
(%)				(100.0)			(100.0)				
腎・尿路腫瘍		1	1				2		2	6.50	3.54
(%)		(50.0)	(50.0)				(100.0)				
閉塞性尿路障害				2			2		2	11.50	2.12
(%)				(100.0)			(100.0)				
骨髄腫											
(%)											
腎形成不全				2			2		2	10.00	
(%)				(100.0)			(100.0)				
不明		9	54	30	5	17	115		115	10.37	5.19
(%)		(7.8)	(47.0)	(26.1)	(4.3)	(14.8)	(100.0)				
再導入		1	7	8	1	1	18	1	19	9.89	4.57
(%)		(5.6)	(38.9)	(44.4)	(5.6)	(5.6)	(100.0)				
その他		4	9	17		2	32		32	9.69	4.13
(%)		(12.5)	(28.1)	(53.1)		(6.3)	(100.0)				
合計		139	618	615	97	131	1,600	17	1,617	9.95	4.81
(%)		(8.7)	(38.6)	(38.4)	(6.1)	(8.2)	(100.0)				
記載なし											
(%)											
総計		139	618	615	97	131	1,600	17	1,617	9.95	4.81
(%)		(8.7)	(38.6)	(38.4)	(6.1)	(8.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査