

表 2114 身長(cm) 治療方法(新分類)別 (透析患者全体)

治療方法 (新分類)	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
施設血液透析	248	567	2,337	8,637	19,227	30,751	35,744	43,719	39,134	24,965	9,096	2,404	386	59	217,274	31,451	248,725	160.03	9.62
(%)	(0.1)	(0.3)	(1.1)	(4.0)	(8.8)	(14.2)	(16.5)	(20.1)	(18.0)	(11.5)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				
Off-line HDF	5	7	40	178	437	718	836	911	793	577	244	71	14	2	4,833	499	5,332	160.21	9.70
(%)	(0.1)	(0.1)	(0.8)	(3.7)	(9.0)	(14.9)	(17.3)	(18.8)	(16.4)	(11.9)	(5.0)	(1.5)	(0.3)	(0.0)	(100.0)				
On-line HDF	31	74	305	1,128	2,966	5,327	6,340	7,690	7,349	5,296	2,061	573	96	12	39,248	5,279	44,527	161.15	9.44
(%)	(0.1)	(0.2)	(0.8)	(2.9)	(7.6)	(13.6)	(16.2)	(19.6)	(18.7)	(13.5)	(5.3)	(1.5)	(0.2)	(0.0)	(100.0)				
Push/Pull HDF	1		1	3	9	22	10	19	15	12	7	1			100	10	110	159.83	10.20
(%)	(1.0)		(1.0)	(3.0)	(9.0)	(22.0)	(10.0)	(19.0)	(15.0)	(12.0)	(7.0)	(1.0)			(100.0)				
バイオフィльтраーション		1	4	15	24	43	30	37	36	29	14				233	34	267	158.89	10.36
(%)		(0.4)	(1.7)	(6.4)	(10.3)	(18.5)	(12.9)	(15.9)	(15.5)	(12.4)	(6.0)				(100.0)				
IHDF	2	11	41	168	312	516	553	656	595	371	108	20	5	2	3,360	180	3,540	159.26	9.55
(%)	(0.1)	(0.3)	(1.2)	(5.0)	(9.3)	(15.4)	(16.5)	(19.5)	(17.7)	(11.0)	(3.2)	(0.6)	(0.1)	(0.1)	(100.0)				
血液濾過					1	2	4	3	1	2					13	3	16	159.23	8.25
(%)					(7.7)	(15.4)	(30.8)	(23.1)	(7.7)	(15.4)					(100.0)				
血液吸着透析 (リクセル等使用)		1	13	47	139	264	277	234	175	101	43	2	1		1,297	79	1,376	158.12	8.73
(%)		(0.1)	(1.0)	(3.6)	(10.7)	(20.4)	(21.4)	(18.0)	(13.5)	(7.8)	(3.3)	(0.2)	(0.1)		(100.0)				
在宅血液透析				3	8	25	41	85	90	108	61	17	4	1	443	112	555	166.96	8.36
(%)				(0.7)	(1.8)	(5.6)	(9.3)	(19.2)	(20.3)	(24.4)	(13.8)	(3.8)	(0.9)	(0.2)	(100.0)				
腹膜透析 (CAPD)	7	4	39	129	292	510	613	704	664	445	190	39	2	1	3,639	1,263	4,902	160.54	9.66
(%)	(0.2)	(0.1)	(1.1)	(3.5)	(8.0)	(14.0)	(16.8)	(19.3)	(18.2)	(12.2)	(5.2)	(1.1)	(0.1)	(0.0)	(100.0)				
腹膜透析 (APD)	55	3	24	63	133	244	320	387	407	306	129	33	4	2	2,110	665	2,775	160.05	14.61
(%)	(2.6)	(0.1)	(1.1)	(3.0)	(6.3)	(11.6)	(15.2)	(18.3)	(19.3)	(14.5)	(6.1)	(1.6)	(0.2)	(0.1)	(100.0)				
腹膜透析 (CCPD)	6	1	2	16	39	93	110	187	186	140	75	17	4	1	877	215	1,092	163.19	10.31
(%)	(0.7)	(0.1)	(0.2)	(1.8)	(4.4)	(10.6)	(12.5)	(21.3)	(21.2)	(16.0)	(8.6)	(1.9)	(0.5)	(0.1)	(100.0)				
合計	355	669	2,806	10,387	23,587	38,515	44,878	54,632	49,445	32,352	12,028	3,177	516	80	273,427	39,790	313,217	160.20	9.66
(%)	(0.1)	(0.2)	(1.0)	(3.8)	(8.6)	(14.1)	(16.4)	(20.0)	(18.1)	(11.8)	(4.4)	(1.2)	(0.2)	(0.0)	(100.0)				
記載なし																			
(%)																			
総計	355	669	2,806	10,387	23,587	38,515	44,878	54,632	49,445	32,352	12,028	3,177	516	80	273,427	39,790	313,217	160.20	9.66
(%)	(0.1)	(0.2)	(1.0)	(3.8)	(8.6)	(14.1)	(16.4)	(20.0)	(18.1)	(11.8)	(4.4)	(1.2)	(0.2)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析 (CCPD) は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査